

1 Tensor of Quantitative Equational Theories

2 **Giorgio Bacci** ✉ 

3 Department of Computer Science, Aalborg University, Aalborg, Denmark

4 **Radu Mardare** ✉

5 Department of Computer & Information Sciences, University of Strathclyde, Glasgow, Scotland

6 **Prakash Panangaden** ✉

7 School of Computer Science, McGill University, Montreal, Canada

8 **Gordon Plotkin** ✉

9 LFCS, School of Informatics, University of Edinburgh, Edinburgh, Scotland

10 — Abstract —

11 We develop a theory for the commutative combination of quantitative effects, their *tensor*, given as a
12 combination of quantitative equational theories that imposes mutual commutation of the operations
13 from each theory. As such, it extends the sum of two theories, which is just their unrestrained
14 combination. Tensors of theories arise in several contexts; in particular, in the semantics of
15 programming languages, the monad transformer for global state is given by a tensor.

16 We show that under certain assumptions on the quantitative theories the free monad that arises
17 from the tensor of two theories is the categorical tensor of the free monads on the theories. As an
18 application, we provide the first algebraic axiomatizations of labelled Markov processes and Markov
19 decision processes. Apart from the intrinsic interest in the axiomatizations, it is pleasing they are
20 obtained compositionally by means of the sum and tensor of simpler quantitative equational theories.

21 **2012 ACM Subject Classification** Theory of computation → Equational logic and rewriting

22 **Keywords and phrases** Quantitative equational theories, Tensor, Monads, Quantitative Effects.

23 **Digital Object Identifier** 10.4230/LIPIcs.CALCO.2021.2

24 **1** Introduction

25 The theory of computational effects began with the work of Moggi [25, 26] seeking a unified
26 category-theoretic account of the semantics of higher-order programming languages. He
27 modelled computational effects (which he called notions of computation) by means of strong
28 monads on a base category with cartesian closed structure. With Cenciarelli [5], he later
29 extended the theory by allowing a compositional treatment of various semantic phenomena
30 such as state, IO, exceptions, resumptions, etc, via the use of monad transformers. This work
31 was followed up by the program of Plotkin and Power [27, 28] on an axiomatic understanding
32 of computational effects as arising from operations and equations via the use of Lawvere
33 theories (see also [14]). In a fundamental contribution [12] together with Hyland they
34 developed a unified modular theory for algebraic effects that supports their combination by
35 taking the *sum* and *tensor* of their Lawvere theories. This allowed them to recover in a more
36 pleasing algebraic structural way many of the monad transformers considered by Moggi.

37 Quantitative equational theories, introduced by Mardare et al. [21], are a logical framework
38 generalising the standard concept of equational logic to account for a concept of approximate
39 equality. They are used to describe quantitative effects as monads on categories of metric
40 spaces. Following the work of Hyland et al. [12], in [1] we developed a theory for the sum of
41 quantitative equational theories, and proved that it corresponds to the categorical sum of
42 quantitative effects as monads. As a major example, we axiomatised Markov processes with
43 discounted probabilistic bisimilarity distance [7] as the sum of two theories: interpolative
44 barycentric algebras (which axiomatises probability distributions with the Kantorovich



© Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon Plotkin;
licensed under Creative Commons License CC-BY 4.0

9th Conference on Algebra and Coalgebra in Computer Science (CALCO 2021).

Editors: Fabio Gadducci and Alexandra Silva; Article No. 2; pp. 2:1–2:29

Leibniz International Proceedings in Informatics



LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

metric [21]) and contractive operators (used to express the transition to the next state).

Whereas the sum of two monads is the simplest combination supporting both given effects with no interactions between them, the tensor additionally requires commutation of these effects over each other. Some of the most important monad transformers have an elegant abstract description using tensor. Specifically, Moggi’s transformers for state, reader, and writer are examples of tensors [12].

In the present paper we extend the work initiated in [1], and develop the theory for the *tensor of quantitative equational theories*. The main contributions are:

1. we prove that the tensor of quantitative theories corresponds to the categorical tensor of their induced quantitative effects as strong monads;
2. we give quantitative axiomatisations to the *quantitative reader* and *writer monads*, from which we obtain analogues of Moggi’s transformers at the level of theories using tensor;
3. we provide the first axiomatization of *labelled Markov processes* and *Markov decision processes* with their discounted bisimilarity metrics.

For the proof of (1) we introduce the concept of *pre-operation of a strong functor*, which we use to conveniently characterise the commutative bialgebras for the monads (which correspond to the Eilenberg-Moore algebras for their tensor). Crucially, this allows us to carry out the technical development directly at the level of quantitative equational theories without passing via a correspondence with metric-enriched Lawvere theories.

The axiomatisations in (3) are two major examples for our compositional theory quantitative effects. Specifically, we obtain the discounted bisimilarity metrics for labelled Markov processes and Markov decision processes with rewards by complementing the axiomatization for Markov processes presented in [1]. We model reactions to action labels by tensoring with the theory of quantitative reading computations (corresponding to Moggi’s reader monad transformer); while rewards are recovered by tensoring with the theory of quantitative writing computations (corresponding to Moggi’s writer monad transformer). We will illustrate our compositional approach by decomposing the proposed axiomatisations into their basic components and showing how to combine them step-by-step to get the desired result.

Further Related Work. In [12, 11] the tensor of (enriched) Lawvere theories is characterized as the colimit of certain commutative cocones, and the correspondence with the tensor of monads is obtained via the equivalence between Lawvere theories and monads. Since it is not hard to show that (basic) quantitative equational theories can be characterised as metric-enriched Lawvere theories, one may think to recover the correspondence with the tensor of monads via the equivalence with Lawvere theories. Alas, quantitative equational theories and Lawvere theories are not equivalent, as the latter allows generic operations with metric spaces as arities, while the framework of Mardare et al. [21] does not. An equivalence with *discrete* Lawvere theories [13] (where arities are just countable ordinals) does not hold either, because quantitative equations implicitly impose the existence operations with non-discrete arities which cannot be expressed in the framework of discrete Lawvere theories.

The above arguments required us to follow a different path, which led us to the introduction of pre-operations for a strong functor F . Pre-operations are related to Plokin and Power’s *algebraic operations* [29, 30] in the sense that their assignment to F -algebras are the appropriate version of algebraic operations for functors. Moreover, when considered over a strong monad T they correspond to generic effects of type $I \rightarrow Tv$ (*i.e.*, Kleisli maps of type $I \rightarrow v$, where I is the identity for the monoidal product). The reason why we consider pre-operations over functors, and not just monads, is to relate the operations of an algebraic monad with those of its signature. This was crucial in the technical development of Section 5.

92 Finally, we remark that quantitative equational theories are a natural kind of enriched
 93 equational theory expressive enough to recover many examples of interest in computer science
 94 (see [21, 1, 24]), but not corresponding to metric-enriched Lawvere theories. In this respect,
 95 it is nice that also this simpler subclass of enriched theories are closed under sum and tensor.

96 2 Preliminaries and Notation

97 An *extended metric space* is a pair (X, d) consisting of a set X equipped with a distance
 98 function $d: X \times X \rightarrow [0, \infty]$ allowed to have infinite values, satisfying: (i) $d(x, y) = 0$ iff
 99 $x = y$, (ii) $d(x, y) = d(y, x)$ and (iii) $d(x, z) \leq d(x, y) + d(y, z)$.

100 A sequence (x_i) in X is *Cauchy* if $\forall \epsilon > 0, \exists N, \forall i, j \geq N, d(x_i, x_j) \leq \epsilon$. If every Cauchy
 101 sequence converges, the extended metric space (X, d) is said to be *complete*. If a space is not
 102 complete it can be completed by a well-known construction called *Cauchy completion*. We
 103 write $\overline{(X, d)}$, or just \overline{X} , for the completion of (X, d) .

104 We denote by **Met** the category of extended metric spaces with morphisms the non-
 105 expansive maps, *i.e.* the $f: (X, d_X) \rightarrow (Y, d_Y)$ such that $d_X(x, y) \geq d_Y(f(x), f(y))$. This
 106 category is both complete (*i.e.*, have all limits) and cocomplete (*i.e.*, have all colimits). We
 107 will consider also the full subcategory **CMet** of complete extended metric spaces.

108 The categorical properties of extended metric spaces are much nicer than usual metric
 109 spaces. In particular, we note that **Met** is a symmetric monoidal category, with monoidal
 110 product $(X, d_X) \square (Y, d_Y)$ being the extended metric space with underlying set $X \times Y$ and
 111 extended metric $d_{X \square Y}((x, y)(x', y')) = d_X(x, x') + d_Y(y, y')$ (*cf.* [19]). Note that this is not
 112 the cartesian product in **Met** (for which $+$ above would be replaced by \max).

113 The monoidal product \square introduced above defines a closed monoidal structure on **Met**,
 114 with internal hom $[(X, d_X), (Y, d_Y)]$ given by the set of non-expansive maps from X to Y
 115 with point-wise supremum extended metric $d_{[X, Y]}(f, g) = \sup_{x \in X} d(f(x), g(x))$.

116 Finally, we recall the basic definitions of strong functor (and monad), strong natural
 117 transformations, and fix the notation (for more details see e.g. [17, 18]). Let \mathbf{V} be a symmetric
 118 monoidal closed category with monoidal product¹ $\square: \mathbf{V} \times \mathbf{V} \rightarrow \mathbf{V}$, unit object $I \in \mathbf{V}$, and
 119 internal hom-functor $[-, -]: \mathbf{V} \times \mathbf{V} \rightarrow \mathbf{V}$. We will denote the *counit* (or evaluation map) of
 120 the adjunction $(V \square -) \dashv [V, -]$ by $ev^V: V \square [V, -] \Rightarrow Id$ and the *unit* (or co-evaluation
 121 map) by $\bar{ev}^V: Id \rightarrow [V, V \square -]$.

122 A functor $F: \mathbf{V} \rightarrow \mathbf{V}$ is *strong* with *monoidal strength* $t_{V, W}: V \square F(W) \rightarrow F(V \square$
 123 $W)$, if t is a natural transformation satisfying the coherence conditions $F\lambda \circ t = \lambda$ and
 124 $t \circ (id \square t) \circ \alpha = F\alpha \circ t$, w.r.t. the associator α and left unitor λ of \mathbf{V} . The dual strength
 125 $\hat{t}_{V, W}: F(W) \square V \rightarrow F(W \square V)$ is given by $\hat{t} = F(s) \circ t \circ s$, where $s: V \square W \rightarrow W \square V$ is
 126 the natural isomorphism of the symmetric monoidal category \mathbf{V} . A natural transformation
 127 $\theta: F \Rightarrow G$ is said *strong* if F, G are strong functors with strengths t, σ , respectively, and
 128 $\sigma \circ (id \square \theta) = \theta \circ t$, meaning that θ interacts well with the strengths.

129 A monad (T, η, μ) with unit $\eta: Id \Rightarrow T$ and multiplication $\mu: TT \Rightarrow T$, is *strong* if T is a
 130 strong functor with strength t such that $t \circ (id \square \eta) = \eta$ and $\mu \circ tt = t \circ (id \square \mu)$.

131 Note that strong functors (resp. monads) on a symmetric monoidal closed category \mathbf{V}
 132 are equivalent to \mathbf{V} -enriched functors (resp. monads) on the self-enriched category \mathbf{V} [17].

¹ We denote the monoidal product by \square to avoid confusion with other tensorial operations we will deal with in this paper, e.g., the tensor of monads.

133 3 Quantitative Equational Theories

134 Quantitative equations were introduced in [21]. In this framework equalities $t \equiv_\varepsilon s$ are
 135 indexed by a positive rational number, to capture the idea that t is “within ε ” of s . This
 136 informal notion is formalised in a manner analogous to traditional equational logic. In this
 137 section we review this formalism.

138 Let Σ be a signature of function symbols $f: n \in \Sigma$ of arity $n \in \mathbb{N}$. Let X be a countable
 139 set of variables, ranged over by x, y, z, \dots . We write $\mathbb{T}(\Sigma, X)$ for the set of Σ -terms freely
 140 generated over X , ranged over by t, s, u, \dots .

141 A *substitution of type Σ* is a function $\sigma: X \rightarrow \mathbb{T}(\Sigma, X)$, canonically extended to terms as
 142 $\sigma(f(t_1, \dots, t_n)) = f(\sigma(t_1), \dots, \sigma(t_n))$; we write $\mathcal{S}(\Sigma)$ for the set of substitutions of type Σ .

143 A *quantitative equation of type Σ over X* is an expression of the form $t \equiv_\varepsilon s$, for
 144 $t, s \in \mathbb{T}(\Sigma, X)$ and $\varepsilon \in \mathbb{Q}_{\geq 0}$. We use $\mathcal{V}(\Sigma, X)$ to denote the set of quantitative equations of
 145 type Σ over X , and its subsets will be ranged over by Γ, Θ, \dots . Let $\mathcal{E}(\Sigma, X)$ be the set of
 146 *conditional quantitative equations* on $\mathbb{T}(\Sigma, X)$, which are expressions of the form

$$147 \quad \{t_1 \equiv_{\varepsilon_1} s_1, \dots, t_n \equiv_{\varepsilon_n} s_n\} \vdash t \equiv_\varepsilon s,$$

148 for arbitrary $s_i, t_i, s, t \in \mathbb{T}(\Sigma, X)$ and $\varepsilon_i, \varepsilon \in \mathbb{Q}_{\geq 0}$.

149 A *quantitative equational theory of type Σ over X* is a set \mathcal{U} of conditional quantitative
 150 equations on $\mathbb{T}(\Sigma, X)$ containing the axioms and closed under the rules of inference below, for
 151 arbitrary $x, y, z, x_i, y_i \in X$, terms $s, t \in \mathbb{T}(\Sigma, X)$, rationals $\varepsilon, \varepsilon' \in \mathbb{Q}_{\geq 0}$, and $\Gamma, \Theta \subseteq \mathcal{V}(\Sigma, X)$,

$$152 \quad (\text{Refl}) \vdash x \equiv_0 x,$$

$$153 \quad (\text{Symm}) \{x \equiv_\varepsilon y\} \vdash y \equiv_\varepsilon x,$$

$$154 \quad (\text{Triang}) \{x \equiv_\varepsilon z, z \equiv_{\varepsilon'} y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y,$$

$$155 \quad (\text{Max}) \{x \equiv_\varepsilon y\} \vdash x \equiv_{\varepsilon+\varepsilon'} y, \text{ for all } \varepsilon' > 0,$$

$$156 \quad (\text{Cont}) \{x \equiv_{\varepsilon'} y \mid \varepsilon' > \varepsilon\} \vdash x \equiv_\varepsilon y,$$

$$157 \quad (f\text{-NE}) \{x_i \equiv_\varepsilon y_i \mid i=1..n\} \vdash f(x_1, \dots, x_n) \equiv_\varepsilon f(y_1, \dots, y_n), \text{ for } f: n \in \Sigma,$$

$$158 \quad (\text{Subst}) \text{ If } \Gamma \vdash t \equiv_\varepsilon s, \text{ then } \{\sigma(t) \equiv_\varepsilon \sigma(s) \mid t \equiv_\varepsilon s \in \Gamma\} \vdash \sigma(t) \equiv_\varepsilon \sigma(s), \text{ for } \sigma \in \mathcal{S}(\Sigma),$$

$$159 \quad (\text{Ass}) \text{ If } t \equiv_\varepsilon s \in \Gamma, \text{ then } \Gamma \vdash t \equiv_\varepsilon s,$$

$$160 \quad (\text{Cut}) \text{ If } \Gamma \vdash \Theta \text{ and } \Theta \vdash t \equiv_\varepsilon s, \text{ then } \Gamma \vdash t \equiv_\varepsilon s,$$

162 where we write $\Gamma \vdash \Theta$ to mean that $\Gamma \vdash t \equiv_\varepsilon s$ holds for all $t \equiv_\varepsilon s \in \Theta$.

163 The rules (Subst), (Cut), (Ass) are the usual rules of equational logic. The axioms
 164 (Refl), (Symm), (Triang) correspond, respectively, to reflexivity, symmetry, and the triangle
 165 inequality; (Max) represents inclusion of neighborhoods of increasing diameter; (Cont) is the
 166 limiting property of a decreasing chain of neighborhoods with converging diameters; and
 167 (f -NE) expresses non-expansiveness of $f \in \Sigma$.

168 A set A of conditional quantitative equations *axiomatizes* a quantitative equational theory
 169 \mathcal{U} , if \mathcal{U} is the smallest quantitative equational theory containing A .

170 The models of these theories, called *quantitative Σ -algebras*, are Σ -algebras in **Met**.

171 ► **Definition 1** (Quantitative Algebra). *A quantitative Σ -algebra is a tuple $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, where
 172 A is an extended metric space and $\Sigma^{\mathcal{A}} = \{f^{\mathcal{A}}: A^n \rightarrow A \mid f: n \in \Sigma\}$ is a set of non-expansive
 173 interpretations (i.e., satisfying $\max_i d_A(a_i, b_i) \geq d_A(f^{\mathcal{A}}(a_1, \dots, a_n), f^{\mathcal{A}}(b_1, \dots, b_n))$).*

174 The morphisms between quantitative Σ -algebras are non-expansive Σ -homomorphisms.
 175 Quantitative Σ -algebras and their morphism form a category, denoted by **QA**(Σ).

176 $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$ satisfies the conditional quantitative equation $\Gamma \vdash t \equiv_{\varepsilon} s$ in $\mathcal{E}(\Sigma, X)$, written
 177 $\Gamma \Vdash_{\mathcal{A}} t \equiv_{\varepsilon} s$, if for any assignment $\iota: X \rightarrow A$, the following implication holds

$$178 \quad (\forall t' \equiv_{\varepsilon'} s' \in \Gamma, d_A(\iota(t'), \iota(s')) \leq \varepsilon') \Rightarrow d_A(\iota(t), \iota(s)) \leq \varepsilon,$$

180 where $\iota(t)$ is the homomorphic interpretation of t in \mathcal{A} .

181 A quantitative algebra \mathcal{A} is said to *satisfy* (or be a *model* for) the quantitative theory \mathcal{U} ,
 182 if $\Gamma \Vdash_{\mathcal{A}} t \equiv_{\varepsilon} s$ whenever $\Gamma \vdash t \equiv_{\varepsilon} s \in \mathcal{U}$. We write $\mathbb{K}(\Sigma, \mathcal{U})$ for the collection of models of a
 183 theory \mathcal{U} of type Σ .

184 Sometimes it is convenient to consider the quantitative Σ -algebras whose carrier is a
 185 complete extended metric space. This class of algebras forms a full subcategory of $\mathbf{QA}(\Sigma)$,
 186 written $\mathbf{CQA}(\Sigma)$. Similarly, we write $\mathbb{CK}(\Sigma, \mathcal{U})$ for the full subcategory of quantitative
 187 Σ -algebras in $\mathbf{CQA}(\Sigma)$ which are models of \mathcal{U} .

188 The following lifts the Cauchy completion of metric spaces to quantitative algebras.

189 **► Definition 2.** (*Algebra Completion*) The Cauchy completion of a quantitative Σ -algebra
 190 $\mathcal{A} = (A, \Sigma^{\mathcal{A}})$, is the quantitative Σ -algebra $\bar{\mathcal{A}} = (\bar{A}, \Sigma^{\bar{\mathcal{A}}})$, where \bar{A} is the Cauchy completion
 191 of A and $\Sigma^{\bar{\mathcal{A}}} = \{f^{\bar{\mathcal{A}}}: \bar{A}^n \rightarrow \bar{A} \mid f: n \in \Sigma\}$ is such that for Cauchy sequences $(b_j^i)_j$ converging
 192 to $b^i \in \bar{A}$, for $1 \leq i \leq n$, $f^{\bar{\mathcal{A}}}(b^1, \dots, b^n) = \lim_j f^{\mathcal{A}}(b_j^1, \dots, b_j^n)$.

193 The above extends to a functor $\mathbb{C}: \mathbf{QA}(\Sigma) \rightarrow \mathbf{CQA}(\Sigma)$ which is the left adjoint to the
 194 functor embedding $\mathbf{CQA}(\Sigma)$ into $\mathbf{QA}(\Sigma)$.

195 The completion of quantitative Σ -algebras extends also to a functor from $\mathbb{K}(\Sigma, \mathcal{U})$ to
 196 $\mathbb{CK}(\Sigma, \mathcal{U})$, whenever \mathcal{U} can be axiomatised by a collection of *continuous schemata*, which are
 197 conditional quantitative equations of the form

$$198 \quad \{x_i \equiv_{\varepsilon_i} y_i \mid i = 1..n\} \vdash t \equiv_{\varepsilon} s, \quad \text{for all } \varepsilon \geq f(\varepsilon_1, \dots, \varepsilon_n),$$

200 where $f: \mathbb{R}_{\geq 0}^n \rightarrow \mathbb{R}_{\geq 0}$ is a continuous real-valued function, $\varepsilon, \varepsilon_i \in \mathbb{Q}_{\geq 0}$, and $x_i, y_i \in X$. We
 201 call such a theory *continuous*.

202 Free Monads on Quantitative Equational Theories

203 To every signature Σ , one can associate a *signature endofunctor* (also called Σ) on \mathbf{Met} by:

$$204 \quad \Sigma = \coprod_{f: n \in \Sigma} Id^n.$$

205 It is easy to see that, by couniversality of the coproduct, quantitative Σ -algebras correspond
 206 to Σ -algebras for the functor Σ in \mathbf{Met} , and the morphisms between them to non-expansive
 207 homomorphisms of Σ -algebras. Below we pass between the two points of view as convenient.

208 **► Theorem 3** (Free Algebra [21]). *The forgetful functor $\mathbb{K}(\Sigma, \mathcal{U}) \rightarrow \mathbf{Met}$ has a left adjoint.*

209 The left adjoint assigns to any $X \in \mathbf{Met}$ a *free quantitative Σ -algebra* $(T_X, \psi_X^{\mathcal{U}})$ satisfying \mathcal{U} ,
 210 from which one canonically obtains the monad $(T_{\mathcal{U}}, \eta^{\mathcal{U}}, \mu^{\mathcal{U}})$, with functor $T_{\mathcal{U}}: \mathbf{Met} \rightarrow \mathbf{Met}$
 211 mapping $X \in \mathbf{Met}$ to the carrier T_X of the free algebra.

212 A similar free construction also holds for quantitative algebras in $\mathbf{CQA}(\Sigma)$ for continuous
 213 quantitative equational theories, implying that the forgetful functor from $\mathbb{CK}(\Sigma, \mathcal{U})$ to \mathbf{CMet}
 214 has a left adjoint. In particular, $\mathbb{CT}_{\mathcal{U}}$ is the free monad on \mathcal{U} in \mathbf{CMet} , provided that the
 215 quantitative equational theory is continuous.

216 Finally, let $T\text{-Alg}$ denote the category of Eilenberg-Moore (EM) algebras for a monad T .
 217 In [1], it is shown that, whenever the quantitative theory \mathcal{U} is *basic*, *i.e.*, it can be axiomatised
 218 by a set of conditional equations of the form

$$219 \quad \{x_1 \equiv_{\varepsilon_1} y_1, \dots, x_n \equiv_{\varepsilon_n} y_n\} \vdash t \equiv_{\varepsilon} s,$$

220 where $x_i, y_i \in X$ (*cf.* [22]), then EM $T_{\mathcal{U}}$ -algebras are in 1-1 correspondence with the quantita-
 221 tive algebras satisfying \mathcal{U} :

222 ► **Theorem 4.** *For any basic quantitative equational theory \mathcal{U} of type Σ , $T_{\mathcal{U}}\text{-Alg} \cong \mathbb{K}(\Sigma, \mathcal{U})$.*

223 4 Tensor of Strong Monads

224 In this section we provide the definition of *tensor of strong monads* on a generic symmetric
 225 monoidal closed category \mathbf{V} . The presentation follows and generalises that of Manes [20],
 226 which considers only the case of monads on \mathbf{Set} .

227 Let v be an object in \mathbf{V} . As \mathbf{V} is self-enriched, it has all v -fold *powers* (or v -*powers*)
 228 X^v , of any object $X \in \mathbf{V}$, defined as $X^v = [v, X]$ [16]. Moreover, $(-)^v: \mathbf{V} \rightarrow \mathbf{V}$ is a strong
 229 functor with strength $\xi_{X,Y}: X \square Y^v \rightarrow (X \square Y)^v$ obtained by currying

$$230 \quad v \square (X \square Y^v) \xrightarrow{\cong} X \square (v \square Y^v) \xrightarrow{X \square ev} X \square Y.$$

231 Let $F: \mathbf{V} \rightarrow \mathbf{V}$ be a strong functor with strength t . The v -*power* functor $(-)^v$ can be
 232 lifted to F -algebras by mapping (A, a) to $(A, a)^v = (A^v, a^v \circ \sigma_A)$, where $\sigma_A: FA^v \Rightarrow (FA)^v$ is
 233 the strong natural transformation obtained from t by currying $Fev_A^v \circ t_{v, A^v}$. Hence F -algebras
 234 are closed under powers of \mathbf{V} -objects.

235 ► **Definition 5** (Pre-operation of a strong functor). *Let $F: \mathbf{V} \rightarrow \mathbf{V}$ be a strong functor and*
 236 *$v \in \mathbf{V}$. A v -ary pre-operation of F is a strong natural transformation of type $(-)^v \Rightarrow F$.*

237 We denote by $\mathcal{O}_F(v)$ the set of v -ary pre-operations of F . An application of $g \in \mathcal{O}_F(v)$
 238 to an F -algebra (A, a) is the composite $a^g = a \circ g_A$. We call a^g an *operation* of (A, a) .

239 ► **Proposition 6.** *Let $(A, a), (B, b)$ be F -algebras of a strong endofunctor F on \mathbf{V} and*
 240 *$f: A \rightarrow B$ a morphism in \mathbf{V} . Then, the following are equivalent:*

- 241 1. *f is an F -homomorphism from (A, a) to (B, b) ;*
- 242 2. *For every $v \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$, $f \circ a^g = b^g \circ f^v$.*

243 The above proposition indicates that F -algebras are precisely characterised by their
 244 operations. In some situations, depending on the functor F , one gets the same characterisation
 245 with much fewer operations. We identify this property with the following definition.

246 ► **Definition 7** (Density). *A set \mathcal{D} of pre-operations of a strong functor $F: \mathbf{V} \rightarrow \mathbf{V}$ is dense,*
 247 *if for any F -algebras $(A, a), (B, b)$ and $f: A \rightarrow B$ in \mathbf{V} , the following are equivalent:*

- 248 1. *f is an F -homomorphism from (A, a) to (B, b) ;*
- 249 2. *For every v -ary pre-operation $g \in \mathcal{D}$, $f \circ a^g = b^g \circ f^v$.*

250 Let F, G be two strong endofunctors on \mathbf{V} . A $\langle F, G \rangle$ -*bialgebra* is a triple (A, a, b) consisting
 251 of an object $A \in \mathbf{V}$ with both an F -algebra structure $a: FA \rightarrow A$ and a G -algebra structure
 252 $b: GA \rightarrow A$. A morphism of $\langle F, G \rangle$ -bialgebras is an arrow that is simultaneously an F - and
 253 G -homomorphism. Denote by $\langle F, G \rangle\text{-biAlg}$ the category of $\langle F, G \rangle$ -bialgebras.

254 ► **Proposition 8.** *Let (A, a, b) be an $\langle F, G \rangle$ -bialgebra. The following statements are equivalent:*

- 255 1. For all $v \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$, a^g is a G -homomorphism;
 256 2. For all $w \in \mathbf{V}$ and $h \in \mathcal{O}_G(w)$, b^h is an F -homomorphism.
 257 Diagrammatically:

$$\begin{array}{ccc}
 GA^v \xrightarrow{\bar{b}} A^v & & FA^w \xrightarrow{\bar{a}} A^w \\
 G(a^g) \downarrow & (1) \quad \downarrow a^g & \text{iff} \quad F(b^h) \downarrow & (2) \quad \downarrow b^h \\
 GA \xrightarrow{b} A & & FA \xrightarrow{a} A
 \end{array}$$

259 where $(A, a)^w = (A^w, \bar{a})$ and $(A, b)^v = (A^v, \bar{b})$.

260 ► **Definition 9** (Commutative bialgebra). A $\langle F, G \rangle$ -bialgebra (A, a, b) is commutative if it
 261 satisfies either of the equivalent conditions of Proposition 8.

262 In the case the functors F and G admit dense sets of pre-operations, commutativity for
 263 their bialgebras can be more conveniently expressed in the following way.

264 ► **Proposition 10.** Let \mathcal{D} and \mathcal{E} be dense sets of pre-operations for F and G , respectively. A
 265 $\langle F, G \rangle$ -bialgebra (A, a, b) is commutative iff it satisfies either of the equivalent conditions:

- 266 1. For all $g \in \mathcal{D}$, a^g is a G -homomorphism;
 267 2. For all $h \in \mathcal{E}$, b^h is an F -homomorphism.

268 Let (T, η, μ) be a strong monad on \mathbf{V} . Note that, as T is a strong functor and the
 269 EM-algebras for T are closed under powers of \mathbf{V} -objects, all the results and definitions given
 270 in this section extends to EM-algebras for T .

271 Let (T, η, μ) , (T', η', μ') be two strong monads on \mathbf{V} . A EM $\langle T, T' \rangle$ -bialgebra is a triple
 272 (A, a, a') consisting of an object $A \in \mathbf{V}$ with both an EM T -algebra structure $a: TA \rightarrow A$
 273 and an EM T' -algebra structure $a': T'A \rightarrow A$. We say that an EM $\langle T, T' \rangle$ -bialgebra
 274 (A, a, a') is *commutative* if it is so as a $\langle T, T' \rangle$ -bialgebra for the functors T, T' . We denote
 275 by $\langle T, T' \rangle$ -**biAlg** the category of EM $\langle T, T' \rangle$ -bialgebras and by $(T \otimes T')$ -**biAlg**, the full
 276 subcategory of the commutative EM $\langle T, T' \rangle$ -bialgebras.

277 ► **Definition 11** (Tensor of monads). If the forgetful functor $(T \otimes T')$ -**biAlg** $\rightarrow \mathbf{V}$ has a left
 278 adjoint, then the monad induced by the adjunction is the tensor of T, T' , denoted $T \otimes T'$.

279 Note that the tensor of monads does not necessarily exist (see [4] for counterexamples).
 280 However, when it does $T \otimes T' \cong T' \otimes T$, as the categories of commutative biagebras
 281 $(T \otimes T')$ -**biAlg** and $(T' \otimes T)$ -**biAlg** are isomorphic.

282 5 Tensor of Quantitative Theories

283 In this section, we develop the theory for the *tensor* of quantitative equational theories. The
 284 main result is that the free monad on the tensor of two theories is the tensor of the monads
 285 on the theories. In the proof given, we use the fact that the quantitative theories are *basic*,
 286 as this allows us to exploit the correspondence between the algebras of a theory \mathcal{U} and the
 287 EM-algebras of the monad $T_{\mathcal{U}}$ (Theorem 4).

288 Let Σ, Σ' be two disjoint signatures. Following Freyd [8] (and [12]), we define the tensor
 289 of two quantitative equational theories $\mathcal{U}, \mathcal{U}'$ of respective types Σ and Σ' , written $\mathcal{U} \otimes \mathcal{U}'$,
 290 as the smallest quantitative theory containing $\mathcal{U}, \mathcal{U}'$ and the quantitative equations

$$291 \vdash f(g(x_1^1, \dots, x_m^1), \dots, g(x_1^n, \dots, x_m^n)) \equiv_0 g(f(x_1^1, \dots, x_1^n), \dots, f(x_m^1, \dots, x_m^n)), \quad (1)$$

292 for all $f: n \in \Sigma$ and $g: m \in \Sigma'$, expressing that the operations of one theory commute with
 293 the operations of the other.

294 **5.1 Density of Symbolic Pre-operations**

 295 Towards our main result, we identify a dense set of pre-operations for the free monads on
 296 quantitative equational theories which, in turn, will gives us a simpler characterization for
 297 commutative bialgebras for these monads (*cf.* Proposition 10).

 298 First notice that any signature functor $\Sigma = \coprod_{f:n \in \Sigma} Id^n$ in **Met** is strong, as it is the
 299 coproduct of the strong functors $Id^n \cong (-)^n$, where $n \in \mathbf{Met}$ denotes the set $\{1, \dots, n\}$
 300 equipped with the discrete extended metric assigning infinite distance to distinct elements.
 301 Moreover, the injections $in_f: (-)^n \Rightarrow \Sigma$ are strong natural transformations, hence they are
 302 n -ary pre-operations of Σ (*cf.* Definition 5).

 303 ► **Proposition 12.** $\mathcal{S}_\Sigma = \{in_f \mid f: n \in \Sigma\}$ is a dense set of pre-operations of Σ .

 304 In the following, the pre-operations in \mathcal{S}_Σ will be called *symbolic*, and to simplify the notation,
 305 for any $f: n \in \Sigma$ and Σ -algebra (A, a) , we write a^f instead of a^{in_f} .

 306 Let \mathcal{U} be a quantitative equational theory of type Σ . Then, also the monad $T_{\mathcal{U}}$ is strong,
 307 with strength $\zeta_{X,Y}: X \square T_{\mathcal{U}}Y \rightarrow T_{\mathcal{U}}(X \square Y)$ obtained by uncurrying the unique map $h_{X,Y}$
 308 that, by Theorem 3, makes the following diagram commute

309
$$\begin{array}{ccccc} Y & \xrightarrow{\eta_Y^{\mathcal{U}}} & T_{\mathcal{U}}Y & \xleftarrow{\psi_Y^{\mathcal{U}}} & \Sigma T_{\mathcal{U}}Y \\ \beta_{X,Y} \curvearrowright & & \downarrow h_{X,Y} & & \downarrow \Sigma h_{X,Y} \\ & & (T_{\mathcal{U}}(X \square Y))^X & \xleftarrow{\psi_{(T_{\mathcal{U}}(X \square Y))^X}^{\mathcal{U}}} & \Sigma(T_{\mathcal{U}}(X \square Y))^X \end{array}$$

 310 where $\beta_{X,Y}$ is the currying of $\eta_{X \square Y}^{\mathcal{U}}: X \square Y \rightarrow T_{\mathcal{U}}(X \square Y)$.

 311 Since a monad is strong iff both its unit and multiplication are strong natural transform-
 312 ations, both $\eta^{\mathcal{U}}, \mu^{\mathcal{U}}$ are strong. Moreover, also $\psi^{\mathcal{U}}: \Sigma T_{\mathcal{U}} \Rightarrow T_{\mathcal{U}}$ is strong.

 313 Thus any pre-operation $g \in \mathcal{O}_\Sigma(v)$ can be tuned into an pre-operation of $T_{\mathcal{U}}$ as the
 314 composite

315
$$(-)^v \xrightarrow{g} \Sigma \xrightarrow{\Sigma \eta^{\mathcal{U}}} \Sigma T_{\mathcal{U}} \xrightarrow{\psi^{\mathcal{U}}} T_{\mathcal{U}}.$$

 316 In particular, when the theory \mathcal{U} is basic, by exploiting Theorem 4, the above transforma-
 317 tion allows us to turn any dense set of pre-operations of Σ into a dense set of pre-operations
 318 of $T_{\mathcal{U}}$.

 319 ► **Proposition 13.** Let \mathcal{U} be a basic quantitative theory of type Σ and \mathcal{D} a dense set of
 320 pre-operations of Σ . Then $\{\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g \mid g \in \mathcal{D}\}$ is a dense set of pre-operations of $T_{\mathcal{U}}$.

 321 By combining Propositions 12 and 13, we have that $\mathcal{S}_{T_{\mathcal{U}}} = \{\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ in_f \mid f: n \in \Sigma\}$ is
 322 a dense set of pre-operations of $T_{\mathcal{U}}$. We call also these pre-operations *symbolic* and we simplify
 323 the notation by writing $a^{(f)}$ instead of $a^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ in_f)}$, for $f: n \in \Sigma$ and $(A, a) \in T_{\mathcal{U}}\text{-Alg}$.

 324 Thus, as an immediate consequence of Propositions 10 and 13, we obtain the following
 325 simpler characterization for commutative $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -bialgebras.

 326 ► **Corollary 14.** Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories respectively of type Σ, Σ' . A
 327 $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -bialgebra (A, a, b) is commutative iff it satisfies either of the equivalent conditions

- 328
1. For all $f: n \in \Sigma$, $a^{(f)}$ is a $T_{\mathcal{U}'}$ -homomorphism;
 2. For all $g: n \in \Sigma', b^{(g)}$ is a $T_{\mathcal{U}}$ -homomorphism.
- 329

5.2 Tensor of Free Monads on Quantitative Theories

Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories respectively of type Σ, Σ' . We show that any model for $\mathcal{U} \otimes \mathcal{U}'$ is a $\langle \mathcal{U} \otimes \mathcal{U}' \rangle$ -bialgebra: an extended metric space A with both a Σ -algebra structure $a: \Sigma A \rightarrow A$ satisfying \mathcal{U} and a Σ' -algebra structure $b: \Sigma' A \rightarrow A$ satisfying \mathcal{U}' and respecting the diagrammatic condition below, for all $f: n \in \Sigma$ and $g: m \in \Sigma'$

$$\begin{array}{ccc}
 A^n & \xrightarrow{a^f} & A \leftarrow b^g & A^m \\
 (b^g)^n \uparrow & & & \uparrow (a^f)^m \\
 (A^m)^n & \xrightarrow[\cong]{\chi} & (A^n)^m &
 \end{array} \tag{2}$$

Formally, we denote by $(\mathcal{U} \otimes \mathcal{U}')\text{-biAlg}$ the category of $\langle \mathcal{U} \otimes \mathcal{U}' \rangle$ -bialgebras, with morphisms the non-expansive homomorphisms preserving both algebraic structures. Then, the following isomorphism of categories holds.

► **Proposition 15.** $\mathbb{K}(\Sigma + \Sigma', \mathcal{U} \otimes \mathcal{U}') \cong (\mathcal{U} \otimes \mathcal{U}')\text{-biAlg}$, for $\mathcal{U}, \mathcal{U}'$ basic quantitative theories.

Moreover, by adapting the isomorphism of Theorem 4 and exploiting the density of symbolic pre-operations (cf. Corollary 14) the following is also true.

► **Proposition 16.** $(\mathcal{U} \otimes \mathcal{U}')\text{-biAlg} \cong (T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$, for $\mathcal{U}, \mathcal{U}'$ basic quantitative theories.

By combining the above two propositions we get the main theorem of this section.

► **Theorem 17.** Let $\mathcal{U}, \mathcal{U}'$ be basic quantitative theories. Then, the monad $T_{\mathcal{U} \otimes \mathcal{U}'}$ in \mathbf{Met} is the tensor of monads $T_{\mathcal{U}} \otimes T_{\mathcal{U}'}$.

Proof. By Propositions 15 and 16 the forgetful functor from $(T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$ to \mathbf{Met} has a left adjoint and the monad generated by this adjunction is isomorphic to $T_{\mathcal{U} \otimes \mathcal{U}'}$. Thus, by definition of tensor of monads, $T_{\mathcal{U} \otimes \mathcal{U}'} \cong T_{\mathcal{U}} \otimes T_{\mathcal{U}'}$. ◀

The above results do not require any specific property of \mathbf{Met} , apart that its morphisms are non-expansive maps. Thus, when the quantitative equational theories are continuous, we can reformulate an alternative version of Theorem 17 which is valid in \mathbf{CMet} .

► **Theorem 18.** Let $\mathcal{U}, \mathcal{U}'$ be continuous quantitative theories. Then, $\mathbb{C}T_{\mathcal{U} \otimes \mathcal{U}'}$ in \mathbf{CMet} is the tensor of monads $\mathbb{C}T_{\mathcal{U}} \otimes \mathbb{C}T_{\mathcal{U}'}$.

6 Quantitative Reader Algebras

Let E be a finite set of input values and fix an enumeration $E = \{e_1, \dots, e_n\}$ for it. The quantitative reader algebras of type E are the algebras for the signature

$$\Sigma_{\mathcal{R}_E} = \{r: |E|\}$$

having only one operator r of arity equal to the number of the input values in E , and satisfying the following axioms

$$(\text{Idem}) \vdash x \equiv_0 r(x, \dots, x),$$

$$(\text{Diag}) \vdash r(x_{1,1}, \dots, x_{n,n}) \equiv_0 r(r(x_{1,1}, \dots, x_{1,n}), \dots, r(x_{n,1}, \dots, x_{n,n})).$$

The quantitative theory induced by the axioms above, written \mathcal{R}_E , is called *quantitative theory of reading computations*.

Intuitively, the term $r(t_1, \dots, t_n)$ can be interpreted as the computation that proceeds as t_i after reading the value e_i from its input. The axiom (**Idem**) says that if we ignore the value of the input the reading of it is not observable; (**Diag**) says that the resulting computation after reading the input is the same no matter how many times we read it.

► **Remark 19.** For the binary case ($|E| = 2$) we can think of r as an *if-then-else* statement $b?(S, T)$ checking for the value of a fixed global Boolean variable b and proceeding as S when $b = \text{true}$, and as T otherwise. In this case, (**Idem**) and (**Diag**) express the standard program equivalences $S \equiv b?(S, S)$ and $b?(S, T) \equiv b?(b?(S, T'), b?(S', T))$.

In the following, when the set E is clear from the context, we use \mathcal{R} in place of \mathcal{R}_E .

On Metric Spaces

Let E be a finite set. We denote by \underline{E} the extended metric space on E equipped with the indiscrete metric that assigns infinite distance to any pair of distinct elements.

Consider the \underline{E} -power functor $(-)^{\underline{E}}: \mathbf{Met} \rightarrow \mathbf{Met}$, assigning to each $X \in \mathbf{Met}$ the metric space $[E, X]$ of (necessarily non-expansive) maps from \underline{E} to X .

This functor has a monad structure, with unit $\kappa: Id \Rightarrow (-)^{\underline{E}}$ and multiplication $\Delta: ((-)^{\underline{E}})^{\underline{E}} \Rightarrow (-)^{\underline{E}}$, respectively given as follows, for $x \in X$, $e \in E$, and $f \in E \rightarrow X^E$

$$\kappa_X(x)(e) = x, \quad \Delta_X(f)(e) = f(e)(e).$$

This is also known as *reader monad* (also called *environment monad* or *function monad*).

► **Remark 20.** The reader monad is always well defined in a cartesian closed category. Fix an object E . The reader monad $(-)^E$ has unit and multiplication respectively given by

$$X \cong X^1 \xrightarrow{X^!} X^E \quad \text{and} \quad (X^E)^E \cong X^{E \times E} \xrightarrow{X^\delta} X^E,$$

where $!: E \rightarrow 1$ is the unique map to the terminal object and $\delta: E \rightarrow E \times E$ the diagonal map $\delta = \langle id, id \rangle$. However, this definition does not generalise to arbitrary monoidal closed categories, and \mathbf{Met} is such a counterexample. The specific problem with \mathbf{Met} is that $\delta: E \rightarrow E \square E$ is not well-defined for arbitrary $E \in \mathbf{Met}$, as non-expansiveness requires that

$$d_E(e, e') \geq d_{E \square E}(\delta(e), \delta(e')) = d_E(e, e') + d_E(e, e'),$$

which holds only when E has the discrete metric. This is the reason why in our treatment we restrict the set of input values to have discrete metric.

The reader monad $(-)^{\underline{E}}$ is isomorphic to the free monad $T_{\mathcal{R}}$. In other words, the quantitative theory \mathcal{R} of reading computations axiomatises the reader monad.

► **Theorem 21.** *The monads $T_{\mathcal{R}}$ and $(-)^{\underline{E}}$ in \mathbf{Met} are isomorphic.*

Let T be a strong monad with strength t . The natural transformation $\lambda_X: TX^{\underline{E}} \Rightarrow (TX)^{\underline{E}}$ obtained from the strength t by currying $Tev_X^{\underline{E}} \circ t_{\underline{E}, X^{\underline{E}}}$, is a distributive law of monads. Distributive laws induce a notion of monad composition [2], so Moggi's reader monad transformer $T \mapsto (T-)^{\underline{E}}$ is also available in \mathbf{Met} . The following says that we can recover this monad transformer as the operation of tensoring with the reader monad.

► **Theorem 22 (Tensoring with Reader Monad).** *Let T be a strong monad. Then, $T \otimes (-)^{\underline{E}}$ exists and is given as the monad composition $(T-)^{\underline{E}}$.*

By using the above result in combination with Theorem 17, we obtain an analogous transformer at the level of quantitative equational theories as follows.

► **Corollary 23.** *Let \mathcal{U} be a basic quantitative equational theory. Then, $(T_{\mathcal{U}}-)^{\underline{E}}$ is the free monad on the theory $\mathcal{U} \otimes \mathcal{R}$ in \mathbf{Met} .*

409 On Complete Metric Spaces

410 The category \mathbf{CMet} has finite products. Since, we assumed the set of input values E to be
 411 finite, the functor $(-)^E$ is isomorphic to the finite product $(-)^n$, for $n = |E|$. Therefore the
 412 power functor $(-)^E$, preserves Cauchy completeness and can be restricted to an endofunctor
 413 on \mathbf{CMet} . Thus also the reader monad restricts to \mathbf{CMet} .

414 Because \mathcal{R} is a continuous quantitative theory, the free monad on \mathcal{R} in \mathbf{CMet} is $\mathbb{C}T_{\mathcal{R}}$.
 415 Thus, by restricting Theorem 21 to quantitative algebras over \mathbf{CMet} , we obtain:

416 ▶ **Theorem 24.** *The monads $\mathbb{C}T_{\mathcal{R}}$ and $(-)^E$ in \mathbf{CMet} are isomorphic.*

417 In virtue of the above characterisation, by instantiating Theorem 22 in the category of
 418 complete extended metric spaces, in combination with Theorems 17 we obtain the following
 419 variant of the quantitative reader theory transformer on continuous quantitative theories.

420 ▶ **Corollary 25.** *Let \mathcal{U} be a continuous quantitative theory. Then, $(\mathbb{C}T_{\mathcal{U}})^E$ is the free
 421 monad on the theory $\mathcal{U} \otimes \mathcal{R}$ in \mathbf{CMet} .*

422 7 Quantitative Writer Algebras

423 Fix an extended metric space $\Lambda \in \mathbf{Met}$ of *output values* having monoid structure $(\Lambda, *, 0)$
 424 with non-expansive multiplication operation $*$: $\Lambda \times \Lambda \rightarrow \Lambda$.

425 The *quantitative writer algebras* of type Λ are the algebras for the signature

$$426 \Sigma_{\mathcal{W}_{\Lambda}} = \{w_{\alpha} : 1 \mid \alpha \in \Lambda\}$$

427 having a unary operator w_{α} , for each output value $\alpha \in \Lambda$, and satisfying the following axioms

$$428 (\text{Zero}) \vdash x \equiv_0 w_0(x),$$

$$429 (\text{Mult}) \vdash w_{\alpha}(w_{\alpha'}(x)) \equiv_0 w_{\alpha * \alpha'}(x),$$

$$430 (\text{Diff}) \{x \equiv_{\varepsilon} x'\} \vdash w_{\alpha}(x) \equiv_{\delta} w_{\alpha'}(x'), \text{ for } \delta \geq d_{\Lambda}(\alpha, \alpha') + \varepsilon.$$

432 The quantitative theory induced by the axioms above, written \mathcal{W}_{Λ} , is called *quantitative
 433 theory of writing computations*.

434 The term $w_{\alpha}(t)$ represents the computation that proceeds as t after writing α on the
 435 output tape. The axiom (Zero) says that writing the identity element 0 is not observable on
 436 the tape; (Mult) says that consecutive writing operations are stored in the tape in the order
 437 of execution; (Diff) compares two computations w.r.t. the distance of their output values.

438 In the following, when the metric space Λ of output values is clear from the context, we
 439 use \mathcal{W} in place of \mathcal{W}_{Λ} .

440 On Metric Spaces

441 Let $(\Lambda \square -): \mathbf{Met} \rightarrow \mathbf{Met}$ be the functor assigning to each extended metric space X the
 442 space $(\Lambda \square X)$. By exploiting the monoid structure of Λ , the functor $(\Lambda \square -)$ can be given a
 443 monad structure with unit $\tau: Id \Rightarrow (\Lambda \square -)$ and multiplication $\varsigma: (\Lambda \square (\Lambda \square -)) \Rightarrow (\Lambda \square -)$,
 444 respectively given as follows, for arbitrary $x \in X$ and $\alpha, \alpha' \in \Lambda$

$$445 \tau_X(x) = (0, x), \quad \varsigma_X((\alpha, (\alpha', x))) = (\alpha * \alpha', x).$$

447 This monad is also known as *writer monad* (also called *complexity monad*). Note that,
 448 the non-expansiveness of the maps above crucially depends on the assumption that the
 449 multiplication $*$ in Λ is non-expansive.

2:12 Tensor of Quantitative Equational Theories

450 The next theorem says that the writer monad $(\Lambda \square -)$ has a quantitative equational
451 presentation in terms of the theory \mathcal{W} of writing computations.

452 ► **Theorem 26.** *The monads $T_{\mathcal{W}}$ and $(\Lambda \square -)$ in \mathbf{Met} are isomorphic.*

453 Let T be a strong monad with strength t . There is a canonical distributive law of the
454 monad $(\Lambda \square -)$ over T , obtained using the strength $t_{\Lambda, -} : (\Lambda \square T-) \Rightarrow T(\Lambda \square -)$ of T . So
455 $T(\Lambda \square -)$ acquires a canonical monad structure [2], and we obtain Moggi's writer monad
456 transformer $T \mapsto T(\Lambda \square -)$ in \mathbf{Met} .

457 In [12], Hyland et al. observed that Moggi's writer monad transformer can be equivalently
458 recovered as the operation of tensoring with the writer monad.

459 ► **Theorem 27** (Tensoring with Writer Monad [12]). *Let T be a strong monad. Then, the
460 monad composition $T(\Lambda \square -)$ is given as $T \otimes (\Lambda \square -)$.*

461 By combining the above with Theorems 17 and 26, we get an analogous transformer at
462 the level of quantitative equational theories as follows:

463 ► **Corollary 28.** *Let \mathcal{U} be a basic quantitative theory. Then, $T_{\mathcal{U}}(\Lambda \square -)$ is the free monad
464 on the theory $\mathcal{U} \otimes \mathcal{W}$ in \mathbf{Met} .*

465 On Complete Metric Spaces

466 If we assume the monoid $(\Lambda, *, 0)$ to be over a complete extended metric space Λ , the writer
467 monad $(\Lambda \square -)$ is well defined also in \mathbf{CMet} .

468 Since \mathcal{W} is axiomatised by a continuous schema of quantitative conditional equations the
469 free monad on \mathcal{W} in \mathbf{CMet} is given by $\mathbf{CT}_{\mathcal{W}}$. Thus, by restricting the use of Theorem 26 to
470 quantitative algebras over complete extended metric spaces, we obtain:

471 ► **Theorem 29.** *The monads $\mathbf{CT}_{\mathcal{W}}$ and $(\Lambda \square -)$ in \mathbf{CMet} are isomorphic.*

472 Thus, by similar arguments as before, we obtain the following variant of Corollary 28.

473 ► **Corollary 30.** *Let \mathcal{U} be a continuous quantitative theory. Then, $\mathbf{CT}_{\mathcal{U}}(\Lambda \square -)$ is the free
474 monad on the theory $\mathcal{U} \otimes \mathcal{W}$ in \mathbf{CMet} .*

475 **8** The Algebras of Labeled Markov Processes

476 In this section we show how to obtain a quantitative equational axiomatization of labelled
477 Markov processes with discounted bisimilarity metric as the combination, via sum and tensor,
478 of the following simpler quantitative equational theories:

479 (a) *The quantitative theory of interpolative barycentric algebras \mathcal{B} from [21] over the signature
480 $\Sigma_{\mathcal{B}} = \{+_e : 2 \mid e \in [0, 1]\}$ extends M. H. Stone's theory of barycentric algebras [32] (a.k.a.
481 abstract convex algebras) with the following axiom*

$$482 \quad (\text{IB}) \quad \{x \equiv_{\varepsilon} y, x' \equiv_{\varepsilon'} y'\} \vdash x +_e x' \equiv_{\delta} y +_e y', \text{ for } \delta \geq e\varepsilon + (1 - e)\varepsilon'$$

484 expressing that the distance between convex combinations is obtained as the convex
485 interpolation of the distance of their sub-terms. This theory will be used to axiomatise
486 probability distributions with Kantorovich metric [15].

487 (b) *The pointed quantitative theory, defined as the free quantitative theory $\mathcal{U}_{\mathbf{0}}$ (i.e., the
488 one imposing no additional axioms) for a signature $\Sigma_{\mathbf{0}} = \{\mathbf{0} : \mathbf{0}\}$ consisting of a single
489 constant $\mathbf{0}$ symbol. This will be used to axiomatise termination.*

490 (c) The quantitative theory \mathcal{R}_A of reading computations (cf. Section 6) will be used to
 491 axiomatise the reaction to the choice of a label from a set A of action labels.

492 (d) The quantitative theory of contractive operators discussed in [1], is the theory obtained
 493 by imposing a Lipschitz contractive axiom for each operator in the signature. In our case,
 494 we consider a signature $\Sigma_\diamond = \{\diamond: 1\}$ with only one unary operator and the contractive
 495 theory \mathcal{U}_\diamond generated from the axiom

$$496 \quad (\diamond\text{-Lip}) \quad \{x =_\varepsilon y\} \vdash \diamond(x) \equiv_\delta \diamond(y), \text{ for } \delta \geq c\varepsilon,$$

498 where $c \in (0, 1)$ is a fixed *contractive factor* for the operator \diamond . This theory will be used
 499 to axiomatise the transition to a next state with discount factor c .

500 Formally, we define the quantitative theory \mathcal{U}_{LMP} of labelled Markov processes as the
 501 following combination of quantitative theories, with signature Σ_{LMP} given by the disjoint
 502 union of those from its component theories:

$$503 \quad \Sigma_{\text{LMP}} = \Sigma_{\mathcal{B}} + \Sigma_{\mathbf{0}} + \Sigma_{\mathcal{R}_A} + \Sigma_\diamond, \quad \mathcal{U}_{\text{LMP}} = ((\mathcal{B} + \mathcal{U}_{\mathbf{0}}) \otimes \mathcal{R}_A) + \mathcal{U}_\diamond.$$

505 Following [33, Section 6], we regard A -labelled Markov processes over extended metric
 506 spaces as $(\Delta(1 + -))^{\mathbb{A}}$ -coalgebras in \mathbf{Met} , where Δ is the *Kantorovich functor* assigning to
 507 each $X \in \mathbf{Met}$ the space of Radon probability measures with finite moment over X equipped
 508 with Kantorovich metric. In [33] it is shown that the *probabilistic bisimilarity distance* on a
 509 labelled Markov processes (X, τ) is equal to the (pseudo)metric

$$510 \quad \mathbf{d}_{(X, \tau)}(x, x') = d_Z(h(x), h(x')),$$

511 where $h: X \rightarrow Z$ is the unique homomorphism to the final coalgebra (Z, ω) .

512 Similarly to [1], we slightly extend the type of the coalgebras to encompass the case when
 513 the probabilistic bisimilarity distance is discounted by a factor $0 < c < 1$. Explicitly, we
 514 consider coalgebras for the functor $(\Delta(1 + c \cdot -))^{\mathbb{A}}$, where $(c \cdot -)$ is the *rescaling functor*,
 515 mapping a metric space (X, d_X) to $(X, c \cdot d_X)$. This will not change the essence of the results
 516 from [33] that are used in this section to characterise the probabilistic bisimilarity metric.

517 In the remainder of the section we prove that the theory \mathcal{U}_{LMP} axiomatizes (the monad
 518 of) A -labelled Markov processes with c -discounted bisimilarity metric.

519 On Metric Spaces

520 We characterise the monad $T_{\mathcal{U}_{\text{LMP}}}$ in steps. First, note that $T_{\mathcal{U}_{\mathbf{0}}} \cong 1^* = (1 + -)$ is the
 521 *maybe monad*, i.e., freely generated monad on the constant terminal object functor 1. As the
 522 monad $(1 + -)$ is isomorphic to $(1F)^*$, for any functor F , by [1, Theorems 4.4 and 5.2], and
 523 [12, Theorem 4], we obtain the following isomorphism of monads in \mathbf{Met} :

$$524 \quad T_{\mathcal{B} + \mathcal{U}_{\mathbf{0}}} \cong T_{\mathcal{B}} + T_{\mathcal{U}_{\mathbf{0}}} \cong \Pi(1 + -),$$

525 where $\Pi(1 + -)$ is the *finite sub-distribution monad* with functor assigning to $X \in \mathbf{Met}$ the
 526 space of finitely supported Borel sub-probability measures with Kantorovich metric. Thus,
 527 $\mathcal{B} + \mathcal{U}_{\mathbf{0}}$ axiomatizes finitely supported sub-probability distributions with Kantorovich metric.

528 From the above, Theorem 17 and Corollary 23, we further get the monad isomorphism

$$529 \quad T_{(\mathcal{B} + \mathcal{U}_{\mathbf{0}}) \otimes \mathcal{R}_A} \cong \Pi(1 + -) \otimes (-)^{\mathbb{A}} \cong (\Pi(1 + -))^{\mathbb{A}},$$

530 saying that tensoring with the theory \mathcal{R}_A of reading computations corresponds to axiomatic-
 531 ally adding the capability of reacting to the choice of an action label.

532 By [1, Theorem 6.3], $T_{\mathcal{U}_\diamond}$ is isomorphic to the free monad over the rescaling functor $(c \cdot -)$.
 533 Hence, by [1, Theorem 4.4] and [12, Corollary 2] we get the following last isomorphism

$$534 \quad T_{\mathcal{U}_{\mathbf{LMP}}} = T_{((\mathcal{B} + \mathcal{U}_\diamond) \otimes \mathcal{R}_A) + \mathcal{U}_\diamond} \cong \mu y. (\Pi(1 + c \cdot y + -))^A.$$

535 Explicitly, this means that, the free monad on $\mathcal{U}_{\mathbf{LMP}}$ assigns to an arbitrary metric space
 536 $X \in \mathbf{Met}$ the *initial solution* of the following functorial equation in \mathbf{Met}

$$537 \quad LMP_X \cong (\Pi(1 + c \cdot LMP_X + X))^A.$$

538 In particular, when $X = 0$ is the empty metric space (*i.e.*, the initial object in \mathbf{Met}) the
 539 above corresponds to the isomorphism on the initial $(\Pi(1 + c \cdot -))^A$ -algebra. The isomorphism
 540 gives us also a $(\Pi(1 + c \cdot -))^A$ -coalgebra structure on LMP_0 , which can be converted into a
 541 labeled Markov process (LMP_0, τ_0) via a post-composition with the inclusion $\Pi(-) \hookrightarrow \Delta(-)$.

542 The key aspect is that the metric of LMP_0 is exactly the bisimilarity metric.

543 ► **Lemma 31.** d_{LMP_0} is the c -discounted probabilistic bisimilarity metric on (LMP_0, τ_0) .

544 ► **Remark 32.** For a less abstract description of (LMP_0, τ_0) , notice that the elements of LMP_0
 545 are (equivalence classes of) ground terms over the signature $\Sigma_{\mathbf{LMP}}$, which one can interpret
 546 as pointed (or rooted) acyclic labelled Markov processes quotiented by bisimilarity.

547 On Complete Metric Spaces

548 Since all the quantitative theories considered are continuous, we can replicate the same steps
 549 also while interpreting the theory $\mathcal{U}_{\mathbf{LMP}}$ over complete metric spaces, obtaining the monad

$$550 \quad \mathbb{C}T_{\mathcal{U}_{\mathbf{LMP}}} \cong \mu y. \Delta(1 + c \cdot y + -)^A.$$

551 By following similar arguments to [1, Section 8.3], one can prove that the functorial
 552 equation $LMP_X \cong \Delta(1 + c \cdot LMP_X + X)^A$ has a unique solution. Thus by applying the
 553 monad above on $X = 0$ we recover the carrier of the final $(\Delta(1 + c \cdot -))^A$ -coalgebra, equipped
 554 with c -discounted probabilistic bisimilarity metric.

555 ► **Remark 33.** While by interpreting the theory $\mathcal{U}_{\mathbf{LMP}}$ over \mathbf{Met} we can only characterise
 556 Markov processes that are acyclic, by doing it over \mathbf{CMet} we obtain an algebraic representa-
 557 tion of all bisimilarity classes as the elements of the final coalgebra. Thus, among others, we
 558 also recover Markov processes with cyclic structures as the limit of all their finite unfoldings.

559 9 The Algebras of Markov Decision Processes with Rewards

560 As a last example, we provide a quantitative axiomatization of Markov decision processes
 561 with rewards equipped with discounted bisimilarity metric. As the construction is similar to
 562 Section 8, we avoid repeating the details of each step of the monad characterization.

563 Let $(\mathbb{R}, +, 0)$ be the standard monoid structure on the reals. We define the quantitative
 564 theory $\mathcal{U}_{\mathbf{MDP}}$ of Markov decision processes with real-valued rewards as follows

$$565 \quad \Sigma_{\mathbf{MDP}} = \Sigma_{\mathcal{B}} + \Sigma_{\mathcal{W}_{\mathbb{R}}} + \Sigma_{\mathcal{R}_A} + \Sigma_{\diamond}, \quad \mathcal{U}_{\mathbf{MDP}} = ((\mathcal{B} \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}) \otimes \mathcal{R}_A) + \mathcal{U}_{\diamond},$$

566 where $\mathcal{W}_{\mathbb{R}}$ is the theory of writing computations and the other theories are as in Section 8.

567 For convenience, we regard Markov decision processes over metric spaces as the coalgebras
 568 for the functor $(\Delta(\mathbb{R} \square c \cdot -))^A$ on \mathbf{Met} , where the endofunctor $(\mathbb{R} \square -)$ is used to encode
 569 the metric differences at each decision step for the real-valued reward available for two states.
 570 Via this coalgebraic representation, the c -discounted *probabilistic bisimilarity distance* on
 571 this structures can be defined as in [33] (following the same definition of Section 8).
 572

573 ▶ Remark 34. In [31] a Markov decision process is defined as a tuple $(S, p(\cdot|s, a), r(s, a))$
 574 with a Markov kernel $p: S \times A \rightarrow \Delta(S)$ and randomised reward function $r: S \times A \rightarrow \Delta(\mathbb{R})$.
 575 Our coalgebraic representation is the natural generalisation over metric spaces, where the
 576 randomness of the Markov kernel and reward function is combined as a probability measure
 577 on $(\mathbb{R} \square c \cdot S)$, by regarding \mathbb{R} and S as extended metric spaces (for each $a \in A$).

578 On Metric Spaces and Complete Metric Spaces

579 Similarly to what we have done in Section 8 for labelled Markov processes, we relate Markov
 580 decision processes and their c -discounted probabilistic bisimilarity pseudometric with the
 581 free monads on the theory \mathcal{U}_{MDP} in **Met** and **CMet**.

582 The only step that changes in the characterisation of $T_{\mathcal{U}_{\text{MDP}}}$, regards the combination of
 583 theories $\mathcal{B} \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}$, which is dealt using Corollary 28. Thus, similarly to Section 8 we get

$$584 \quad T_{\mathcal{U}_{\text{MDP}}} = T_{((\mathcal{B} + \otimes \mathcal{U}_{\mathcal{W}_{\mathbb{R}}}) \otimes \mathcal{R}_A) + \mathcal{U}_{\diamond}} \cong \mu y. \Pi((\mathbb{R} \square y) + -)^A.$$

585 The metric on the initial solution for the functorial fixed point definition corresponds to the
 586 c -discounted probabilistic bisimilarity (pseudo)metric on its coalgebra structure.

587 Similar considerations apply also when interpreting the theories in the category **CMet** of
 588 complete metric spaces, as the argument follows without issues because \mathbb{R} a complete metric
 589 space. Thus we obtain the following characterisation for the monad:

$$590 \quad \mathbb{C}T_{\mathcal{U}_{\text{LMP}}} \cong \mu y. \Delta((\mathbb{R} \square y) + -)^A.$$

591 Again, the metric on the solution for the above functorial fixed point definition corresponds
 592 to the c -discounted probabilistic bisimilarity metric. Moreover, as the fixed point solution is
 593 unique, $\mathbb{C}T_{\mathcal{U}_{\text{LMP}}}0$ is an algebraic characterization of the final $(\Delta(\mathbb{R} \square c \cdot -))^A$ -coalgebra.

594 10 Conclusions

595 We studied the commutative combination of quantitative effects as the tensor of their
 596 quantitative equational theories. The key result in this regard is Theorem 17, asserting
 597 that the tensor of two quantitative theories corresponds to the categorical tensor of their
 598 free monads. In addition to this general result, we show how to extend to the quantitative
 599 algebraic setting Moggi's notions of reader and writer monad transformers.

600 We illustrate the applicability of our theoretical development with two examples: labeled
 601 Markov processes and Markov decision processes. Apart from the intrinsic interest in their
 602 quantitative equational axiomatisations, what is particularly pleasant is the systematic
 603 compositional way with which one can obtain quantitative axiomatisations of different
 604 variants of Markov processes by just combining theories as new basic ingredients.

605 An example that escapes our compositional treatment via sum and tensor is the com-
 606 bination of probabilities and non-determinism as illustrated in [24]. A possible future work
 607 in this direction is to extend the combination of theories with another operator: the dis-
 608 tributive tensor (see [13, Section 6]). Following a similar intuition by Cheng [6], we claim
 609 that these correspond in a suitable way to Garner's weak distributive law [9]. Our claim
 610 seems promising in the light of the work [10, 3] which consider equational axiomatisations
 611 combining probabilities and non-determinism.

612 — References —

- 613 1 Giorgio Bacci, Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. An algebraic
614 theory of markov processes. In *LICS*, pages 679–688. ACM, 2018.
- 615 2 Jon Beck. Distributive laws. In *Seminar on Triples and Categorical Homology Theory*, volume
616 80 of Lect. Notes Math., pages 119–140. Springer, 1966.
- 617 3 Filippo Bonchi and Alessio Santamaria. Combining semilattices and semimodules. *CoRR*,
618 abs/2012.14778, 2020.
- 619 4 Nathan J. Bowler, Sergey Goncharov, Paul Blain Levy, and Lutz Schröder. Exploring the
620 boundaries of monad tensorability on set. *Log. Methods Comput. Sci.*, 9(3), 2013.
- 621 5 Pietro Cenciarelli and Eugenio Moggi. A syntactic approach to modularity in denotational
622 semantics. Technical report, CWI, 1993. Proc. 5th. Biennial Meeting on Category Theory and
623 Computer Science.
- 624 6 Eugenia Cheng. Distributive laws for lawvere theories. *Compositionality*, 2:1,
625 May 2020. URL: [http://dx.doi.org/10.32408/](http://dx.doi.org/10.32408/compositionality-2-1)
626 [compositionality-2-1](http://dx.doi.org/10.32408/compositionality-2-1), doi:10.32408/
[compositionality-2-1](http://dx.doi.org/10.32408/compositionality-2-1).
- 627 7 Josee Desharnais, Vineet Gupta, Radha Jagadeesan, and Prakash Panangaden. Metrics for
628 labelled Markov processes. *Theoretical Computer Science*, 318(3):323–354, 2004.
- 629 8 Peter J. Freyd. Algebra valued functors in general and tensor products in particular. *Colloq.*
630 *Math.*, 14:89–106, 1966.
- 631 9 Richard Garner. The vitoris monad and weak distributive laws. *Appl. Categorical Struct.*,
632 28(2):339–354, 2020.
- 633 10 Alexandre Goy and Daniela Petrisan. Combining probabilistic and non-deterministic choice
634 via weak distributive laws. In *LICS*, pages 454–464. ACM, 2020.
- 635 11 Martin Hyland, Paul Blain Levy, Gordon D. Plotkin, and John Power. Combining algebraic
636 effects with continuations. *Theor. Comput. Sci.*, 375(1-3):20–40, 2007.
- 637 12 Martin Hyland, Gordon D. Plotkin, and John Power. Combining effects: Sum and tensor.
638 *Theor. Comput. Sci.*, 357(1-3):70–99, 2006. doi:10.1016/j.tcs.2006.03.013.
- 639 13 Martin Hyland and John Power. Discrete lawvere theories and computational effects. *Theor.*
640 *Comput. Sci.*, 366(1-2):144–162, 2006.
- 641 14 Martin Hyland and John Power. The category theoretic understanding of universal algebra:
642 Lawvere theories and monads. *Electronic Notes in Theor. Comp. Sci.*, 172:437–458, 2007.
- 643 15 Leonid Vitalevich Kantorovich. On the transfer of masses (in Russian). *Doklady Akademii*
644 *Nauk*, 5(5-6):1–4, 1942. Translated in *Management Science*, 1958.
- 645 16 Gregory M. Kelly. Basic concepts of enriched category theory. *Theory and Applications of*
646 *Categories*, 1982. Reprinted in 2005.
- 647 17 Anders Kock. Strong functors and monoidal monads. *Arch. Math. (Basel)*, 23:113–120, 1972.
- 648 18 Saunders Mac Lane. *Categories for the Working Mathematician*. Graduate Texts in Mathem-
649 atics. Springer New York, 2nd edition, 1998.
- 650 19 William F. Lawvere. Metric spaces, generalized logic, and closed categories. In *Seminario*
651 *Mat. e. Fis. di Milano*, volume 43, pages 135–166. Springer, 1973.
- 652 20 Ernest Manes. A Triple Theoretic Construction of Compact Algebras. In *Seminar on Triples*
653 *and Categorical Homology Theory*, volume 80 of Lect. Notes Math., pages 91–118. Springer,
654 1966.
- 655 21 Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Quantitative Algebraic Reason-
656 ing. In *LICS*, pages 700–709. ACM, 2016. doi:10.1145/2933575.2934518.
- 657 22 Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. On the axiomatizability
658 of quantitative algebras. In *LICS 2017*, pages 1–12. IEEE Computer Society, 2017. doi:
659 10.1109/LICS.2017.8005102.
- 660 23 Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. Free complete wasserstein
661 algebras. *Log. Methods Comput. Sci.*, 14(3), 2018.

- 662 **24** Matteo Mio and Valeria Vignudelli. Monads and quantitative equational theories for non-
 663 determinism and probability. In *CONCUR*, volume 171 of *LIPIcs*, pages 28:1–28:18. Schloss
 664 Dagstuhl - Leibniz-Zentrum für Informatik, 2020.
- 665 **25** Eugenio Moggi. *The partial lambda calculus*. PhD thesis, University of Edinburgh. College of
 666 Science and Engineering. School of Informatics., 1988.
- 667 **26** Eugenio Moggi. Notions of computation and monads. *Information and computation*, 93(1):55–
 668 92, 1991.
- 669 **27** Gordon Plotkin and John Power. Semantics for algebraic operations. *Electronic Notes in
 670 Theoretical Computer Science*, 45:332–345, 2001.
- 671 **28** Gordon Plotkin and John Power. Notions of computation determine monads. In *Foundations
 672 of Software Science and Computation Structures*, pages 342–356. Springer, 2002.
- 673 **29** Gordon D. Plotkin and John Power. Semantics for algebraic operations. In *MFPS*, volume 45
 674 of *Electronic Notes in Theoretical Computer Science*, pages 332–345. Elsevier, 2001.
- 675 **30** Gordon D. Plotkin and John Power. Algebraic operations and generic effects. *Appl. Categorical
 676 Struct.*, 11(1):69–94, 2003.
- 677 **31** M. L. Puterman. *Markov Decision Processes*. Wiley, 2005.
- 678 **32** Marshall H. Stone. Postulates for the barycentric calculus. *Annali di Matematica Pura ed
 679 Applicata*, 29(1):25–30, 1949.
- 680 **33** Franck van Breugel, Claudio Hermida, Michael Makkai, and James Worrell. Recursively
 681 defined metric spaces without contraction. *Theor. Comput. Sci.*, 380(1-2):143–163, 2007.
 682 doi:10.1016/j.tcs.2007.02.059.

683 **A** Kantorovich Metric on Extended Metric Spaces

684 We assume the reader is familiar with the notions of σ -algebras, measurable functions, and
 685 Borel probability measures. Next we review some facts about metrics between extended
 686 spaces of probability distributions from [23].

687 Let M be an extended metric space. A Borel probability measure μ over M is *Radon* if
 688 for any Borel set $E \subseteq M$, $\mu(E)$ is the supremum of $\mu(K)$ over all compact subsets K of E .
 689 Examples of Radon probability measures are finitely supported probability measures on any
 690 metric space and generic Borel probability measures over complete separable metric spaces.

691 A Radon probability measure μ over M has *finite moment* (of order 1) if, for some
 692 (equivalently all) m_0 , the integral

$$693 \int d_M(m_0, -) d\mu$$

694 is finite. By restricting our attention to Radon measures of finite moment the following is a
 695 well-defined metric [23].

696 The *Kantorovich metric* (or W_1 metric) between Radon probability measures μ, ν of finite
 697 moment over M is defined as:

$$698 \mathcal{K}(d_M)(\mu, \nu) = \min \left\{ \int d_M d\omega \mid \omega \in \mathcal{C}(\mu, \nu) \right\} .$$

699 where $\mathcal{C}(\mu, \nu)$ is the set of *couplings* for a pair of Radon measures (μ, ν) , that is, a Radon
 700 probability measures ω on the product space $M \times M$, such that, for all Borel sets $E \subseteq M$

$$701 \omega(E \times M) = \mu(E) \quad \text{and} \quad \omega(M \times E) = \nu(E) .$$

702
 703 We write $\Delta(M)$ for the set of Radon probability measures with finite moment over M ,
 704 equipped with the Kantorovich metric and $\Pi(M)$ for the subspace of $\Delta(M)$ of the finitely
 705 supported Borel probability measures over M .

706 ► **Theorem 35** (Theorem 2.7 [23]). *Let M be a complete extended metric space. Then, $\Delta(M)$*
 707 *is a complete metric space generated as the Cauchy completion of $\Pi(M)$, i.e., $\overline{\Pi(M)} \cong \Delta(M)$.*

708 **B Interpolative Barycentric Algebras**

709 In this section we recall interpolative barycentric algebras from [21], which are the quantitative
 710 algebras for the signature

$$711 \quad \Sigma_{\mathcal{B}} = \{+_e : 2 \mid e \in [0, 1]\}$$

712 having a binary operator $+_e$, for each $e \in [0, 1]$ (a.k.a. *barycentric signature*), and satisfying
 713 the following axioms

$$714 \quad (\text{B1}) \vdash x +_1 y \equiv_0 x,$$

$$715 \quad (\text{B2}) \vdash x +_e x \equiv_0 x,$$

$$716 \quad (\text{SC}) \vdash x +_e y \equiv_0 y +_{\bar{e}} x,$$

$$717 \quad (\text{SA}) \vdash (x +_e y) +_{e'} z \equiv_0 x +_{ee'} (y +_{\frac{e'-ee'}{1-ee'}} z), \text{ for } e, e' \in [0, 1],$$

$$718 \quad (\text{IB}) \{x \equiv_{\varepsilon} y, x' \equiv_{\varepsilon'} y'\} \vdash x +_e x' \equiv_{\delta} y +_e y', \text{ for } \delta \geq e\varepsilon + (1-e)\varepsilon'.$$

720 The quantitative theory induced by the axioms above, written \mathcal{B} , is called *interpolative*
 721 *barycentric quantitative equational theory*. The axioms (B1), (B2), (SC), (SA) are those of
 722 *barycentric algebras* (a.k.a. *abstract convex sets*) due to M. H. Stone [32] where (SC) stands
 723 for *skew commutativity* and (SA) for *skew associativity*; (IB) is the *interpolative barycentric*
 724 *axiom* introduced in [21].

725 **On Metric Spaces**

726 Let $\Pi: \mathbf{Met} \rightarrow \mathbf{Met}$ be the functor assigning to each $X \in \mathbf{Met}$ the metric space $\Pi(X)$
 727 of finitely supported Borel probability measures with Kantorovich metric and acting on
 728 morphisms $f: X \rightarrow Y$ as $\Pi(f)(\mu) = \mu \circ f^{-1}$, for $\mu \in \Pi(X)$.

729 This functor has a monad structure, with unit $\delta: Id \Rightarrow \Pi$ and multiplication $m: \Pi^2 \Rightarrow \Pi$,
 730 given as follows, for $x \in X$, $\Phi \in \Pi^2(X)$, and Borel subset $E \subseteq X$

$$731 \quad \delta_X(x) = \delta_x, \quad m_X(\Phi)(E) = \int v_E d\Phi,$$

733 where δ_x is the Dirac distribution at x , and $v_E: \Pi(X) \rightarrow [0, 1]$ is the evaluation function,
 734 taking $\mu \in \Pi(X)$ to $\mu(E) \in [0, 1]$. This monad is also known as the *finite distribution monad*.

735 For any $X \in \mathbf{Met}$, one can define a quantitative $\Sigma_{\mathcal{B}}$ -algebra $(\Pi(X), \phi_X)$ as follows, for
 736 arbitrary $\mu, \nu \in \Pi X$

$$737 \quad \phi_X: \Sigma_{\mathcal{B}} \Pi X \rightarrow \Pi X \quad \phi_X(in_{+e}(\mu, \nu)) = e\mu + (1-e)\nu.$$

739 This quantitative algebra satisfies the interpolative barycentric theory \mathcal{B} (cf. [21, The-
 740 orem 10.4]) and is isomorphic to the free quantitative \mathcal{B} -algebra (cf. [21, Theorem 10.5]).

741 Thus, as shown in [21], Π is isomorphic to the free monad $T_{\mathcal{B}}$ on the theory \mathcal{B} of
 742 interpolative barycentric algebras.

743 ► **Theorem 36.** *The monads $T_{\mathcal{B}}$ and Π in \mathbf{Met} are isomorphic.*

744 **On Complete Metric Spaces**

745 Define the functor $\Delta: \mathbf{CMet} \rightarrow \mathbf{CMet}$ assigning to each $X \in \mathbf{CMet}$ the complete metric
 746 space $\Delta(X)$ of Radon probability measures with finite moment and equipped with Kantorovich
 747 metric; acting on morphisms $f: X \rightarrow Y$ as $\Delta(f)(\mu) = \mu \circ f^{-1}$, for $\mu \in \Delta(X)$. This functor
 748 has a monad structure, defined similarly to the one for Π . It is known as the *Kantorovich*
 749 *monad*.

750 By exploiting Lemma 35, one can verify that $\Delta(X)$ with its canonical barycentric algebra
 751 structure is the free interpolative barycentric algebra in \mathbf{CMet} (Theorem 3.7 [23]). As the
 752 canonical monad structure on $\mathbb{C}\Pi$ is isomorphic to the one on Δ in \mathbf{CMet} , by Theorem 36,
 753 we obtain the following.

754 **► Theorem 37.** *The monads $\mathbb{C}T_{\mathcal{B}}$ and Δ in \mathbf{CMet} are isomorphic.*

755 Note that, since \mathcal{B} is axiomatised by a continuous schema of quantitative equations, the
 756 free monad on \mathcal{B} in \mathbf{CMet} is given by $\mathbb{C}T_{\mathcal{B}}$. In other words, [23, Theorem 3.7] provides an
 757 algebraic characterisation of the Kantorovich monad.

758 **C Omitted Proofs**

759 **Proof.** (of Proposition 6) (1) \Rightarrow (2) follows by definition of a^g , b^g and naturality of g .
 760 As for (2) \Rightarrow (1), note that since \mathbf{V} is a symmetric monoidal closed category, we have a
 761 1-1 correspondence between strong and \mathbf{V} -enriched endofunctors on \mathbf{V} , and also between
 762 strong and \mathbf{V} -enriched natural transformations [17]. Therefore, by (the weak form of) the
 763 enriched Yoneda lemma (*cf.* [16]), there exists a natural bijection between strong natural
 764 transformations $g \in \mathcal{O}_F(A)$ and the (generalised) elements of FA , *i.e.*, morphisms of the
 765 form $I \rightarrow FA$, obtained via the composition

$$766 \quad I \xrightarrow{id_A} A^A \xrightarrow{g_A} FA.$$

767 Thus, for any $e: I \rightarrow FA$, there exists $\hat{e} \in \mathcal{O}_F(A)$ such that $\hat{e}_A \circ id_A = e$. Therefore, by
 768 naturality of \hat{e} , definition of $a^{\hat{e}}$, $b^{\hat{e}}$, and (2), the following diagram commute

$$769 \quad \begin{array}{ccccc} I & \xrightarrow{e} & FA & \xrightarrow{a} & A \\ \downarrow e & \searrow \hat{e}_A & \uparrow id_A & \searrow f^A & \downarrow f \\ & & A^A & \xrightarrow{f^A} & B^A \\ & \swarrow \hat{e}_A & & \downarrow \hat{e}_B & \\ FA & \xrightarrow{Ff} & FB & \xrightarrow{b} & B \end{array}$$

770 implying that $f^I \circ a^I = b^I \circ (Ff^I)$. Then (1) follows by the naturality of the isomorphism
 771 $V \xrightarrow{\cong} V^I$ (obtained by currying $\lambda: I \square V \xrightarrow{\cong} V$) and the commutativity of the diagram

$$772 \quad \begin{array}{ccccc} & \xrightarrow{a} & & & \\ \overbrace{FA} & \xrightarrow{\cong} & (FA)^I & \xrightarrow{a^I} & A^I \xleftarrow{\cong} A \\ \downarrow Ff & & \downarrow (Ff)^I & & \downarrow f^I \\ \overbrace{FB} & \xrightarrow{\cong} & (FB)^I & \xrightarrow{b^I} & B^I \xleftarrow{\cong} B \\ & \xrightarrow{b} & & & \end{array}$$

773 ► **Proposition 38.** Let (A, a) be a F -algebra of a strong endofunctor F on \mathbf{V} . Then, for any
 774 $v, w \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$ the following commute

$$775 \begin{array}{ccc} (A^v)^w & \xrightarrow[\cong]{\chi} & (A^w)^v \\ & \searrow^{(a^g)^w} & \swarrow_{\bar{a}^g} \\ & & A^w \end{array}$$

776 where $(A, a)^w = (A^w, \bar{a})$ and χ is the canonical isomorphism.

777 **Proof.** By the universality of the counit $ev: (w \square -) \Rightarrow Id$ of the adjunction $(w \square -) \dashv (-)^w$
 778 it suffices to show that the following two diagrams commute:

$$779 \begin{array}{ccc} w \square (A^v)^w & & w \square (A^v)^w \xrightarrow{w \square \chi} w \square (A^w)^v \\ \swarrow^{a^g \circ ev} \downarrow w \square (a^g)^w & & \downarrow a^g \circ ev \downarrow w \square \bar{a}^g \\ A \xleftarrow{ev} w \square A^w & & A \xleftarrow{ev} w \square A^w \end{array}$$

780 The diagram to the left commutes by naturality of the counit ev ; the one to the right
 781 commutes as follows, where ξ and t are respectively the strengths of $(-)^v$ and F

$$782 \begin{array}{ccccc} & & w \square (A^w)^v & & \\ & \nearrow^{w \square \chi} & \downarrow \xi & \searrow^{w \square g} & \\ w \square (A^v)^w & & (w \square A^w)^v & & \\ \downarrow ev & \swarrow^{ev^v} & \downarrow g & & \\ A^v & & F(w \square A^w) \xleftarrow{t} w \square F A^w & & \\ \downarrow g & \swarrow^{F ev} & \downarrow w \square \sigma & & \\ F A & & w \square (F A)^w & & \\ \downarrow a & \xleftarrow{ev} & \downarrow w \square a^w & & \\ A & \xleftarrow{ev} & w \square A^w & & \end{array}$$

783 by naturality of the counit ev ; definition of ξ and χ ; definition of the law $\sigma: F(-)^w \Rightarrow (F-)^w$;
 784 definition of a^g, \bar{a}^g ; by $\bar{a} = a^w \circ \sigma$; and because g is strong. ◀

785 **Proof.** (of Proposition 8) (1) \Rightarrow (2) By Proposition 6, we prove (2) by showing that for all
 786 $v \in \mathbf{V}$ and $g \in \mathcal{O}_F(v)$, $b^h \circ \bar{a}^g = a^g \circ (b^h)^v$. This is shown by the diagram below

$$787 \begin{array}{ccc} (A^w)^v & \xrightarrow{\bar{a}^g} & A^w \\ \downarrow (b^h)^v \swarrow \chi & & \swarrow (a^g)^w \\ & (A^v)^w & \downarrow b^h \\ & \swarrow \bar{b}^h & \\ A^v & \xrightarrow{a^g} & A \end{array}$$

788 which commutes by Proposition 38, (1), definition of a^g , and naturality of g . The implication
 789 (2) \Rightarrow (1) is similar. ◀

790 **Proof.** (of Proposition 10) The equivalence of the statements (1), (2) follows as in Proposi-
 791 tion 8, by using the density of \mathcal{D} and \mathcal{E} in lieu of Proposition 6.

792 Assume (A, a, b) is a commutative $\langle F, G' \rangle$ -bialgebra. Then, (1) follows trivially because,
 793 \mathcal{D} is a subset of pre-operations of F . For the converse implication, assume (1) and let
 794 $h \in \mathcal{O}_G(w)$ for some $w \in \mathbf{V}$. We want to show that

$$795 \begin{array}{ccc} FA^w & \xrightarrow{\bar{a}} & A^w \\ F(b^h) \downarrow & & \downarrow b^h \\ \Sigma A & \xrightarrow{a} & A \end{array}$$

796 commutes, where $(A, a)^w = (A^w, \bar{a})$. By density of \mathcal{D} , it suffices to show that for all v -ary
 797 pre-operation $g \in \mathcal{D}$, $b^h \circ \bar{a}^g = a^g \circ (b^h)^v$. This follows by

$$798 \begin{array}{ccc} (A^w)^v & \xrightarrow{\bar{a}^g} & A^w \\ \downarrow (b^h)^v & \swarrow \chi & \downarrow b^h \\ & (A^v)^w & \\ \downarrow (b^h)^v & \swarrow \chi^{-1} & \downarrow b^h \\ A^v & \xrightarrow{a^g} & A \end{array}$$

799 which commutes by Proposition 38, (1), definition of a^g , and naturality of g . ◀

800 **Proof.** (of Proposition 12) Let $(A, a), (B, b)$ be Σ -algebras in \mathbf{Met} and $h: A \rightarrow B$ a non-
 801 expansive map. We want to prove the equivalence of

- 802 1. f is a Σ -homomorphisms from (A, a) to (B, b) ;
- 803 2. For every $f: n \in \Sigma$, $h \circ a^f = b^f \circ h^v$.

804 (1) \Rightarrow (2) follows by definition of a^f, b^f and naturality of $in_f: (-)^n \Rightarrow \Sigma$. The implication
 805 (2) \Rightarrow (1) follows by the universality of the coproduct, as $\Sigma = \coprod_{f:n \in \Sigma} Id^n$. ◀

806 ▶ **Proposition 39.** $T_{\mathcal{U}}$ is a strong monad with strength ζ .

807 **Proof.** Naturality of ζ follows by definition and naturality of $\eta^{\mathcal{U}}$ and $\psi^{\mathcal{U}}$. The coherence
 808 conditions of a monoidal strength follow by universality of the evaluation and co-evaluation
 809 maps of the closed structure of \mathbf{Met} , Theorem 3 and definition of $(T_{\mathcal{U}}, \eta^{\mathcal{U}}, \mu^{\mathcal{U}})$. ◀

810 **Proof.** (of Proposition 13) $(A, a), (B, b)$ be $T_{\mathcal{U}}$ -algebras and $h: A \rightarrow B$ a non-expansive map.
 811 We want to prove the equivalence of

- 812 1. h is a $T_{\mathcal{U}}$ -homomorphism from (A, a) to (B, b) ;
- 813 2. For every v -ary pre-operation $g \in \mathcal{D}$, $h \circ a^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)} = b^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)} \circ h^v$.

814 (1) \Rightarrow (2) follows by definition of $a^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)}, b^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)}$ and naturality of $\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g$.
 815 For the converse implication, recall that the isomorphism of categories from Theorem 4, maps
 816 a $T_{\mathcal{U}}$ -algebra (A, a) to the Σ -algebra $(A, a \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})$ and morphisms essentially to themselves.
 817 Thus (2) \Rightarrow (1) follows by density of \mathcal{D} and definition of $a^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)}, b^{(\psi^{\mathcal{U}} \circ \Sigma \eta^{\mathcal{U}} \circ g)}$. ◀

818 **Proof.** (of Proposition 15) The isomorphism is given by the pair of functors

$$819 \mathbb{K}(\Sigma + \Sigma', \mathcal{U} \otimes \mathcal{U}') \begin{array}{c} \xrightarrow{H} \\ \xleftarrow{K} \end{array} \mathbb{K}((\Sigma, \mathcal{U}) \otimes (\Sigma', \mathcal{U}'))$$

820 defined, for a $(\Sigma + \Sigma')$ -algebra (A, a) satisfying $\mathcal{U} \otimes \mathcal{U}'$ and a $\langle \mathcal{U} \otimes \mathcal{U}' \rangle$ -bialgebra (B, b, b') ,
 821 respectively as

$$822 \quad H(A, a) = (A, a \circ in_l, a \circ in_r), \quad K(B, b, b') = (B, [b, b']),$$

824 where $[b, b'] : \Sigma B + \Sigma' B \rightarrow B$ is the unique map induced by b and b' by universality of the
 825 coproduct. Both functors are identity on morphisms; it is easy to see that a homomorphism
 826 in one sense is also a homomorphism in the other.

827 The pair of functors above is the restriction of the isomorphic pair of functors used in the
 828 proof of [1, Proposition 4.1]. Thus, to show H and K are well defined we are just left to deal
 829 with checking that the restriction conditions on the subcategories are preserved both ways.

830 As for H , we prove that whenever $\mathcal{A} = (A, a)$ satisfies the quantitative equation in (1),
 831 then $(A, a \circ in_l, a \circ in_r)$ satisfies the commutativity of the diagram in (2). This follows as,
 832 for all $f : n \in \Sigma$ and $g : m \in \Sigma'$, by definition of algebraic interpretation $(-)^{\mathcal{A}}$, we have

$$833 \quad f^{\mathcal{A}} = a \circ in_l \circ in_f = (a \circ in_l)^f,$$

$$834 \quad g^{\mathcal{A}} = a \circ in_r \circ in_g = (a \circ in_r)^g.$$

836 Thus, the satisfiability (1) coincides with the commutativity of the diagram in (2).

837 For K we need to show that whenever (B, b, b') satisfies the commutativity of the diagram
 838 in (2), then $\mathcal{A} = (A, [b, b'])$ satisfies (1). This follows as, for all $f : n \in \Sigma$ and $g : m \in \Sigma'$, by
 839 definition of algebraic interpretation $(-)^{\mathcal{A}}$, we have

$$840 \quad f^{\mathcal{A}} = [b, b'] \circ in_l \circ in_f = (b)^f,$$

$$841 \quad g^{\mathcal{A}} = [b, b'] \circ in_r \circ in_g = (b')^g.$$

843 Thus, the commutativity of the diagram in (2) coincides with the satisfiability of (1). \blacktriangleleft

844 **Proof.** (of Proposition 16) Recall the isomorphism of categories from Theorem 4

$$845 \quad T_{\mathcal{U}}\text{-Alg} \begin{array}{c} \xrightarrow{H} \\ \xleftarrow{K} \end{array} \mathbb{K}(\Sigma, \mathcal{U})$$

846 mapping morphisms essentially to themselves and on objects acting as follows: for $(A, a) \in$
 847 $T_{\mathcal{U}}\text{-Alg}$ and $(B, b) \in \mathbb{K}(\Sigma, \mathcal{U})$,

$$848 \quad H(A, a) = (A, a \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}}), \quad K(B, b) = (B, b_b),$$

850 where $\beta_b : T_{\mathcal{U}} B \rightarrow B$ is the unique map that, by Theorem 3, satisfies the equations $b_b \circ \eta_B^{\mathcal{U}} =$
 851 id_B and $b_b \circ \psi_B^{\mathcal{U}} = b \circ \Sigma b_b$. (for the details on the proof cf. [1, Theorem 4.2]).

852 Next we show that the obvious point-wise extension of the above functors on the categories
 853 of bialgebras $(\mathcal{U} \otimes \mathcal{U}')\text{-biAlg}$ and $(T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$ is an isomorphism of categories.

854 Clearly, since H and K are inverse with each other, so are their point-wise extensions.
 855 The only thing we are left to prove is that they are well defined; in particular that the
 856 respective commutative conditions are preserved.

857 Let $(A, a, b) \in (T_{\mathcal{U}} \otimes T_{\mathcal{U}'})\text{-biAlg}$. We need to show that condition (2) is satisfied by
 858 $(A, a \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}}, b \circ \psi_A^{\mathcal{U}'} \circ \Sigma \eta_A^{\mathcal{U}'})$. Let $(A, b)^n = (A^n, \bar{b})$. By Corollary 14 and Propositions 12, 13,
 859 we have that the bottom square diagram below commutes for all $f : n \in \Sigma$ and all $g : m \in \Sigma'$,
 860 while the top commute by Proposition 38:

$$861 \quad \begin{array}{ccc} (A^m)^n & & (b^{(g)})^n \\ \downarrow \chi & \searrow & \downarrow \\ (A^n)^m & \xrightarrow{\bar{b}^{(g)}} & A^n \\ (a^{(f)})^m \downarrow & & \downarrow a^{(f)} \\ A^m & \xrightarrow{b^{(g)}} & A \end{array}$$

862 Since $a^{(f)} = (a \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^f$ and $b^{(g)} = (b \circ \psi_A^{\mathcal{U}} \circ \Sigma \eta_A^{\mathcal{U}})^g$, the above diagram proves that
 863 condition (2) holds.

864 Let $(A, a, b) \in (\mathcal{U} \otimes \mathcal{U}')\text{-biAlg}$. We need to show that (A, a_b, b_b) is a $\langle T_{\mathcal{U}}, T_{\mathcal{U}'} \rangle$ -bialgebra.
 865 By Corollary 14, it is sufficient to prove that the following diagram commutes for all $g: m \in \Sigma'$,
 866

$$867 \begin{array}{ccc} T_{\mathcal{U}} A^m & \xrightarrow{\bar{a}_b} & A^m \\ T_{\mathcal{U}}(b_b^{(g)}) \downarrow & & \downarrow b_b^{(g)} \\ T_{\mathcal{U}} A & \xrightarrow{a_b} & A \end{array} \quad (3)$$

868 where $(A, a_b)^m = (A^m, \bar{a}_b)$.

869 Toward proving (3), first notice that the diagram below commutes for all $f: n \in \Sigma$ and
 870 $g: m \in \Sigma'$

$$871 \begin{array}{ccc} (A^n)^m & \xrightarrow{\bar{a}^f} & A^m \\ \downarrow (b^g)^n & \swarrow \chi^{-1} & \downarrow b^g \\ (A^m)^n & \xrightarrow{(a^f)^m} & A^m \\ \downarrow (b^g)^n & \searrow \chi & \downarrow b^g \\ A^n & \xrightarrow{a^f} & A \end{array} \quad (4)$$

872 for $(A, a)^m = (A^m, \bar{a})$ and $(A, b)^n = (A^n, \bar{b})$. Indeed, the bottom commutes because (A, a, b)
 873 satisfies (2), and the top triangle does by Proposition 38.

874 Going back to proving (3), by Theorem 3, it suffices to show that both $b_b^{(g)} \circ \bar{a}_b$ and
 875 $a_b \circ T_{\mathcal{U}}(b_b^{(g)})$ are the (unique) homomorphic extension of a along $b_b^{(g)}$. This is shown by the
 876 following diagrams

$$877 \begin{array}{ccc} A^m & \xrightarrow{\eta^{\mathcal{U}}} & T_{\mathcal{U}} A^m \\ b_b^{(g)} \downarrow & & \downarrow T_{\mathcal{U}} b_b^{(g)} \\ A & \xrightarrow{\eta^{\mathcal{U}}} & T_{\mathcal{U}} A \\ & \searrow id & \downarrow a_b \\ & & A \end{array} \quad \begin{array}{ccc} A^m & \xrightarrow{\eta^{\mathcal{U}}} & T_{\mathcal{U}} A^m \\ & \searrow id & \downarrow \bar{a}_b \\ & & A^m \\ b_b^{(g)} \downarrow & & \downarrow b_b^{(g)} \\ & & A \end{array}$$

$$878 \begin{array}{ccc} T_{\mathcal{U}} A^m & \xleftarrow{\psi^{\mathcal{U}}} & \Sigma T_{\mathcal{U}} A^m \\ T_{\mathcal{U}} b_b^{(g)} \downarrow & & \downarrow \Sigma T_{\mathcal{U}} b_b^{(g)} \\ T_{\mathcal{U}} A & \xleftarrow{\psi^{\mathcal{U}}} & \Sigma T_{\mathcal{U}} A \\ a_b \downarrow & & \downarrow \Sigma a_b \\ A & \xleftarrow{a} & \Sigma A \end{array}$$

$$881 \begin{array}{ccccc} & & T_{\mathcal{U}} A^m & \xleftarrow{\psi^{\mathcal{U}}} & \Sigma T_{\mathcal{U}} A^m \\ & & \bar{a}_b \downarrow & & \downarrow \Sigma \bar{a}_b \\ \Sigma' T_{\mathcal{U}'} A & \xleftarrow{\Sigma' \eta^{\mathcal{U}'}} & A^m & \xleftarrow{\bar{a}} & \Sigma A^m & \xrightarrow{\Sigma \Sigma' \eta^{\mathcal{U}'}} & \Sigma \Sigma' T_{\mathcal{U}'} A \\ \downarrow id & \swarrow & \downarrow in_g & \swarrow \Sigma in_g & \downarrow \Sigma in_g & \searrow & \downarrow id \\ \Sigma' T_{\mathcal{U}'} A & \xleftarrow{\Sigma' b_b} & \Sigma' A & \xrightarrow{\Sigma \Sigma' b_b} & \Sigma \Sigma' A & \xrightarrow{\Sigma \Sigma' b_b} & \Sigma \Sigma' T_{\mathcal{U}'} A \\ \downarrow \psi^{\mathcal{U}'} & & \downarrow b & & \downarrow \Sigma b & & \downarrow \psi^{\mathcal{U}'} \\ T_{\mathcal{U}'} A & \xrightarrow{b_b} & A & \xleftarrow{a} & \Sigma A & \xleftarrow{\Sigma b_b} & \Sigma T_{\mathcal{U}'} \end{array}$$

882 that commute by definitions of $a_b, b_b, b_b^{(g)}, b^g$; by naturality of $\eta^{\mathcal{U}}, \psi^{\mathcal{U}}$; since $(A, a_b)^m$ is a
 883 EM $T_{\mathcal{U}}$ -algebra; because by Theorem 4 $(A, a_b)^m = (A, a)^m$; and since from Propositions 12,
 884 13 and (4) we have that b^g is a Σ -homomorphism. \blacktriangleleft

885 **Proof.** (of Theorem 18) The tensor $\mathcal{U} \otimes \mathcal{U}'$ of continuous theories is also continuous, so that,
 886 by [1, Theorem 3.4], the free monad on it in \mathbf{CMet} is $\mathbf{CT}_{\mathcal{U} \otimes \mathcal{U}'}$. Moreover, by exploiting the
 887 universal property of [1, Theorem 3.4], we can refactor the proofs of Propositions 15 and 16
 888 to obtain the isomorphism $\mathbb{CK}(\Sigma + \Sigma', \mathcal{U} \otimes \mathcal{U}') \cong (\mathbf{CT}_{\mathcal{U}} \otimes \mathbf{CT}_{\mathcal{U}'})\text{-biAlg}$. Thus, by definition
 889 of tensor of monads, $\mathbf{CT}_{\mathcal{U} \otimes \mathcal{U}'} \cong \mathbf{CT}_{\mathcal{U}} \otimes \mathbf{CT}_{\mathcal{U}'}$. \blacktriangleleft

890 C.1 Quantitative Reader Algebras

891 For any $X \in \mathbf{Met}$, we define the quantitative $\Sigma_{\mathcal{R}}$ -algebra (X^E, ρ_X) as follows, for arbitrary
 892 maps $f_1, \dots, f_n: E \rightarrow X$

$$893 \quad \rho_X: \Sigma_{\mathcal{R}} X^E \rightarrow X^E \qquad \rho_X(\text{in}_r(f_1, \dots, f_n))(e_i) = f_i(e_i).$$

895 This quantitative algebra satisfies the quantitative theory \mathcal{R} of reading computations.

896 \blacktriangleright **Proposition 40.** $(X^E, \rho_X) \in \mathbb{K}(\Sigma_{\mathcal{R}}, \mathcal{R})$.

897 **Proof.** Let $r^\rho = \rho_X \circ \text{in}_r$ denote the interpretation of the operator symbol $r: n \in \Sigma_{\mathcal{R}}$ in the
 898 algebra (X^E, ρ_X) . Soundness for the axiom of non-expansiveness (r-NE) follows by the fact
 899 that ρ_X is a well defined map in \mathbf{Met} as shown below

$$900 \quad d_{X^E}(r^\rho(f_1, \dots, f_n), r^\rho(g_1, \dots, g_n))$$

$$901 \quad = \sup_{e_i} d_X(r^\rho(f_1, \dots, f_n)(e_i), r^\rho(g_1, \dots, g_n)(e_i))$$

$$902 \quad = \sup_{e_i} d_X(f_i(e_i), g_i(e_i))$$

$$903 \quad \leq \max_j \left(\sup_{e_i \in E} d_X(f_j(e_i), g_j(e_i)) \right)$$

$$904 \quad \leq \max_j d_{X^E}(f_j, g_j).$$

906 We are left to show that the algebra (X^E, ρ_X) satisfies the axioms (Idem) and (Diag).
 907 Soundness for (Idem) follows by definition of ρ as, for all $e_i \in E$

$$908 \quad r^\rho(f, \dots, f)(e_i) = f(e_i).$$

909 Soundness for (Diag) also follows by definition, as

$$910 \quad r^\rho(r^\rho(f_{1,1}, \dots, f_{1,n}), \dots, r^\rho(f_{n,1}, \dots, f_{n,n}))(e_i)$$

$$911 \quad = r^\rho(f_{i,1}, \dots, f_{i,n})(e_i)$$

$$912 \quad = f_{i,i}(e_i)$$

$$913 \quad = r^\rho(f_{1,1}, \dots, f_{n,n})(e_i).$$

915 Moreover, it is universal in the following sense:

916 \blacktriangleright **Theorem 41.** For any $\Sigma_{\mathcal{R}}$ -algebra (A, a) satisfying \mathcal{R} and non-expansive map $\beta: X \rightarrow A$,
 917 there exists a unique homomorphism $h: X^E \rightarrow A$ of quantitative $\Sigma_{\mathcal{R}}$ -algebras making the
 918 diagram below commute

$$919 \quad \begin{array}{ccccc} X & \xrightarrow{\kappa_X} & X^E & \xleftarrow{\rho_X} & \Sigma_{\mathcal{R}} X^E \\ & \searrow \beta & \downarrow h & & \downarrow \Sigma_{\mathcal{R}} h \\ & & A & \xleftarrow{a} & \Sigma_{\mathcal{R}} A \end{array}$$

920 **Proof.** Let (A, a) be a quantitative $\Sigma_{\mathcal{R}}$ -algebra satisfying \mathcal{R} and $\beta: X \rightarrow A$ a non-expansive
921 map. We define $h: X^E \rightarrow A$ as follows, for arbitrary $f: E \rightarrow X$

$$922 \quad h(f) = a(\text{in}_r(\beta(f(e_1)), \dots, \beta(f(e_n)))) .$$

923 As it is defined as the composition of non-expansive maps, then also h is non-expansive. Next
924 we prove the commutativity of the diagram, that is, $h \circ \kappa_X = \beta$ and $h \circ \rho_X = a \circ \Sigma_{\mathcal{R}} h$.

925 Let $r^\rho = \rho_X \circ \text{in}_r$ and $r^a = a \circ \text{in}_r$ denote the interpretations of $r: n \in \Sigma_{\mathcal{R}}$ in the algebras
926 (X^E, ρ_X) and (A, a) , respectively. Let $x \in X$. Then

$$\begin{aligned} 927 \quad & (h \circ \kappa_X)(x) \\ 928 \quad &= r^a(\beta(\kappa_X(x)(e_1)), \dots, \beta(\kappa_X(x)(e_n))) && \text{(def. } h) \\ 929 \quad &= r^a(\beta(x), \dots, \beta(x)) && \text{(def. } \kappa) \\ 930 \quad &= \beta(x) . && \text{(Idem)} \end{aligned}$$

932 Let $f_1, \dots, f_n: E \rightarrow X$. Then

$$\begin{aligned} 933 \quad & (h \circ \rho_X)(\text{in}_r(f_1, \dots, f_n)) \\ 934 \quad &= r^a(\beta(f_1(e_1)), \dots, \beta(f_n(e_n))) && \text{(def. } h \text{ and } \rho) \\ 935 \quad &= r^a\left(r^a(\beta(f_1(e_1)), \dots, \beta(f_1(e_n))), \dots \right. \\ 936 \quad & \quad \left. \dots, r^a(\beta(f_n(e_1)), \dots, \beta(f_n(e_1)))\right) && \text{(Diag)} \\ 937 \quad &= r^a(h(f_1), \dots, h(f_n)) && \text{(def. } h) \\ 938 \quad &= (a \circ \Sigma_{\mathcal{R}} h)(\text{in}_r(f_1, \dots, f_n)) . && \text{(def. } r^a \text{ and } \Sigma_{\mathcal{R}}) \end{aligned}$$

940 Hence h is a $\Sigma_{\mathcal{R}}$ -homomorphism.

941 It remains to prove the uniqueness of such a homomorphism. Assume there exists
942 $g: X^E \rightarrow A$ such that $g \circ \kappa_X = \beta$ and $g \circ \rho_X = a \circ \Sigma_{\mathcal{R}} g$. Next we prove $h = g$. Notice first
943 that for any $f: X^E \rightarrow X$, $f = r^\rho(\kappa_X(f(e_1)), \dots, \kappa_X(f(e_n)))$, as for all $e_i \in E$, the following
944 holds:

$$\begin{aligned} 945 \quad & f(e_i) = \kappa_X(f(e_i))(e_i) && \text{(def. } \kappa) \\ 946 \quad &= r^\rho(\kappa_X(f(e_1)), \dots, \kappa_X(f(e_n)))(e_i) . && \text{(def. } \rho) \end{aligned}$$

948 From the above we have that, for all $f: X^E \rightarrow X$,

$$\begin{aligned} 949 \quad & h(f) = h(r^\rho(\kappa_X(f(e_1)), \dots, \kappa_X(f(e_n)))) \\ 950 \quad &= r^a((h \circ \kappa)(f(e_1)), \dots, (h \circ \kappa)(f(e_1))) && (h \text{ homo}) \\ 951 \quad &= r^a(\beta(f(e_1)), \dots, \beta(f(e_1))) && (h \circ \kappa = \beta) \\ 952 \quad &= r^a((g \circ \kappa)(f(e_1)), \dots, (g \circ \kappa)(f(e_1))) && (g \circ \kappa = \beta) \\ 953 \quad &= g(r^\rho(\kappa_X(f(e_1)), \dots, \kappa_X(f(e_n)))) && (g \text{ homo}) \\ 954 \quad &= g(f) \\ 955 \end{aligned}$$

956 Therefore, $g = h$. ◀

957 **Proof.** (of Theorem 21) By Theorem 41, the functors $(-)^E$ and $T_{\mathcal{R}}$ are isomorphic and the
958 units of the two monads coincide (up-to iso). We are left to prove that also the multiplications

959 coincide (up-to iso). By Theorem 41, this follows by showing that the following diagram
 960 commutes

$$\begin{array}{ccccc}
 X^E & \xrightarrow{\kappa_{X^E}} & (X^E)^E & \xleftarrow{\rho_{X^E}} & \Sigma_{\mathcal{R}}(X^E)^E \\
 & \searrow id & \downarrow \Delta_X & & \downarrow \Sigma_{\mathcal{R}} \Delta_X \\
 & & X^E & \xleftarrow{\rho_X} & \Sigma_{\mathcal{R}} X^E
 \end{array}$$

962 $\Delta_X \circ \kappa_X = id$ holds since $(-)^E$ is a monad. Finally, the right square diagram commutes as
 963 shown below

$$\begin{aligned}
 964 \quad & (\Delta_X \circ \rho_{X^E})(in_r(F_1, \dots, F_n))(e_i) \\
 965 \quad & = \rho_{X^E}(in_r(F_1, \dots, F_n))(e_i)(e_i) && \text{(def. } \Delta) \\
 966 \quad & = F_i(e_i)(e_i) && \text{(def. } \rho) \\
 967 \quad & = \Delta_X(F_i)(e_i) && \text{(def. } \Delta_X) \\
 968 \quad & = \rho_X(in_r(\Delta_X(F_1), \dots, \Delta_X(F_n)))(e_i) && \text{(def. } \rho) \\
 969 \quad & = (\rho_X \circ \Sigma_{\mathcal{R}} \Delta_X)(in_r(F_1, \dots, F_n))(e_i) && \text{(def. } \Sigma_{\mathcal{R}}) \\
 970
 \end{aligned}$$

971 for arbitrary $F_1, \dots, F_n: E \rightarrow X^E$. ◀

972 **Proof.** (of Theorem 22) Recall that the composite monad $(T-)^E$ is the monad that arises
 973 from the adjunction with the forgetful functor $\lambda\text{-biAlg} \rightarrow \mathbf{Met}$, where $\lambda\text{-biAlg}$ denotes the
 974 full subcategory of EM $\langle T, (-)^E \rangle$ -bialgebras (A, a, b) satisfying the commutativity of the
 975 diagram

$$\begin{array}{ccc}
 TA & \xrightarrow{a} & A \xleftarrow{b} & A^E \\
 Tb \uparrow & & & \uparrow (a)^E \\
 T(A^E) & \xrightarrow{\lambda} & (TA)^E &
 \end{array} \tag{5}$$

977 The bialgebras satisfying (5) are called, λ -bialgebras for the law $\lambda: T(-)^E \Rightarrow (T-)^E$ (see e.g.,
 978 [2]). We show that the category of λ -bialgebras is identical to the category of commutative
 979 $\langle T \otimes (-)^E \rangle$ -bialgebras, that is, that the commutativity of the diagram above corresponds to
 980 either one of the equivalent conditions from Proposition 8.

981 One direction is easy, as if we assume (A, a, b) to be a commutative $\langle T \otimes (-)^E \rangle$ -bialgebra,
 982 then (5) is just the instantiation of (2) from Proposition 8 for $h = id \in \mathcal{O}_{(-)^E}(E)$ as, by
 983 definition of lifting, $(A, a)^E = (A^E, (a)^E \circ \lambda_A)$.

984 For the converse direction, assume (5) holds and let $g \in \mathcal{O}_T(v)$, for some $v \in \mathbf{Met}$.
 985 Then, asking that a^g is a $(-)^E$ -homomorphism (i.e., condition (1) from Proposition 8)
 986 corresponds to the commutativity of the following diagram, as $(A, b)^v = (A^v, b^v \circ \sigma_A)$ and
 987 $(A, a)^E = (A^E, (a)^E \circ \lambda_A)$:

$$\begin{array}{ccccc}
 (A^v)^E & \xrightarrow{\sigma} & (A^E)^v & \xrightarrow{b^v} & A^v \\
 g^E \downarrow & & g \downarrow & & g \downarrow \\
 (TA)^E & & T(A^E) & \xrightarrow{Tb} & TA \\
 & \searrow a^E & \downarrow (a)^E \circ \lambda & & \downarrow a \\
 & & A^E & \xrightarrow{b} & A
 \end{array}$$

989 The bottom-left square is (5), so commutes by hypothesis; the top-right square commutes by
 990 naturality of g ; and finally, the left square commutes by Proposition 38 as, by definitions of
 991 the strengths of $(-)^v$ and $(-)^E$, $\sigma: (A^v)^E \Rightarrow (A^E)^v$ coincides with the canonical isomorphism
 992 (denoted as χ in Proposition 38).

993 Therefore, as the two categories of bialgebras coincide, by definition of tensor of monads,
 994 $T \otimes (-)^E = (T-)^E$. ◀

995 C.2 Quantitative Writer Algebras

996 For any $X \in \mathbf{Met}$, we define the quantitative $\Sigma_{\mathcal{W}}$ -algebra $(\Lambda \square X, \omega_X)$ as follows, for arbitrary
 997 $\alpha, \alpha' \in \Lambda$ and $x \in X$

$$998 \quad \omega_X: \Sigma_{\mathcal{W}}(\Lambda \square X) \rightarrow \Lambda \square X, \quad \omega_X(in_{w_\alpha}(\alpha', x)) = (\alpha * \alpha', x).$$

1000 This quantitative algebra satisfies the quantitative theory \mathcal{W} of writing computations.

1001 ▶ **Proposition 42.** $((\Lambda \square X), \omega_X) \in \mathbb{K}(\Sigma_{\mathcal{W}}, \mathcal{W})$.

1002 **Proof.** Let $w_\alpha^\omega = \omega_X \circ in_{w_\alpha}$ denote the interpretation of the operation $w_\alpha: 1 \in \Sigma_{\mathcal{W}}$ in the
 1003 algebra $(\Lambda \square X, \omega_X)$. Proving the soundness for $(w_\alpha\text{-NE})$, for each $\alpha \in \Lambda$, is equivalent to
 1004 show that the map ω is well-defined in \mathbf{Met} . This follows as

$$\begin{aligned} 1005 \quad & d_{(\Lambda \square X)}(w_\alpha^\omega(\beta, x), w_\alpha^\omega(\beta', x')) \\ 1006 \quad &= d_{(\Lambda \square X)}((\alpha * \beta, x), (\alpha * \beta', x')) && \text{(def. } \omega) \\ 1007 \quad &= d_\Lambda(\alpha * \beta, \alpha * \beta') + d_X(x, x') && \text{(def. } \square) \\ 1008 \quad &\leq \max \left\{ d_\Lambda(\alpha, \alpha), d_\Lambda(\beta, \beta') \right\} + d_X(x, x') && (* \text{ non-exp}) \\ 1009 \quad &= d_\Lambda(\beta, \beta') + d_X(x, x') && \text{(metric)} \\ 1010 \quad &= d_{(\Lambda \square X)}((\beta, x), (\beta', x')). && \text{(def. } \square) \end{aligned}$$

1012 We are missing to prove that the algebra $((\Lambda \square X), \omega_X)$ satisfies the axioms **(Zero)**, **(Mult)**,
 1013 and **(Diff)**. The first one holds trivially as $(\alpha, x) = (0 * \alpha, x)$ because 0 is the identity element
 1014 of the monoid Λ of output values. The soundness of **(Mult)** follows by definition of ω as

$$1015 \quad w_\alpha^\omega(w_{\alpha'}^\omega(\beta, x)) = w_\alpha^\omega((\alpha' * \beta, x)).$$

1016 Finally, soundness for **(Diff)** follows by

$$\begin{aligned} 1017 \quad & d_{(\Lambda \square X)}(w_\alpha^\omega(\beta, x), w_{\alpha'}^\omega(\beta', x')) \\ 1018 \quad &= d_\Lambda(\alpha * \beta, \alpha' * \beta') + d_X(x, x') && \text{(def. } \omega \text{ \& } \square) \\ 1019 \quad &= d_\Lambda(\alpha * \beta, \alpha * \beta') + d_\Lambda(\alpha * \beta', \alpha' * \beta') + d_X(x, x') && \text{(triang. ineq.)} \\ 1020 \quad &\leq d_\Lambda(\beta, \beta') + d_\Lambda(\alpha, \alpha') + d_X(x, x') && (* \text{ non-exp}) \\ 1021 \quad &\geq d_\Lambda(\alpha, \alpha') + d_{(\Lambda \square X)}((\beta, x), (\beta', x')), && \text{(def. } \square) \end{aligned}$$

1023 which concludes our proof. ◀

1024 Moreover, the next result says that this algebra is actually the free quantitative $\Sigma_{\mathcal{W}}$ -
 1025 algebra on X in $\mathbb{K}(\Sigma_{\mathcal{W}}, \mathcal{W})$.

1026 ► **Theorem 43.** For any $\Sigma_{\mathcal{W}}$ -algebra (A, a) satisfying \mathcal{W} and non-expansive map $\beta: X \rightarrow A$,
 1027 there exists a unique homomorphism $h: X^E \rightarrow A$ of quantitative $\Sigma_{\mathcal{W}}$ -algebras making the
 1028 diagram below commute

$$\begin{array}{ccccc}
 X & \xrightarrow{\tau_X} & \Lambda \square X & \xleftarrow{\omega_X} & \Sigma_{\mathcal{W}}(\Lambda \square X) \\
 & \searrow \beta & \downarrow h & & \downarrow \Sigma_{\mathcal{W}} h \\
 & & A & \xleftarrow{a} & \Sigma_{\mathcal{W}} A
 \end{array}$$

1030 **Proof.** Let (A, a) be a $\Sigma_{\mathcal{W}}$ -algebra satisfying \mathcal{W} and $\beta: X \rightarrow A$ a non-expansive map, We
 1031 define the map $h: \Lambda \square X \rightarrow A$ as follows, for arbitrary $\alpha \in \Lambda$ and $x \in X$

$$1032 \quad h((\alpha, x)) = a(in_{w_\alpha}(\beta(x))).$$

1033 Non-expansiveness of h follows by the fact that (A, a) satisfies the axiom (Diff) as shown
 1034 below, where $w_\alpha^a = a \circ in_{w_\alpha}$ denotes the interpretation of $w_\alpha: 1 \in \Sigma_{\mathcal{W}}$ in (A, a) ,

$$\begin{aligned}
 1035 \quad & d_A(h((\alpha, x)), h((\alpha', x'))) \\
 1036 \quad &= d_A(w_\alpha^a(\beta(x)), w_{\alpha'}^a(\beta(x'))) && \text{(def. } h) \\
 1037 \quad &\leq d_\Lambda(\alpha, \alpha') + d_A(\beta(x), \beta(x')) && \text{(Diff)} \\
 1038 \quad &\leq d_\Lambda(\alpha, \alpha') + d_X(x, x') && (\beta \text{ non-exp)} \\
 1039 \quad &= d_{\Lambda \square X}((\alpha, x), (\alpha', x')). && \text{(def. } \square)
 \end{aligned}$$

1041 Next we prove $h \circ \tau_X = \beta$ and $h \circ \omega_X = a \circ \Sigma_{\mathcal{W}} h$.

1042 Let $x \in X$. Then,

$$\begin{aligned}
 1043 \quad & (h \circ \tau_X)(x) = h((0, x)) && \text{(def. } \tau) \\
 1044 \quad &= w_0^a(\beta(x)) && \text{(def. } h) \\
 1045 \quad &= \beta(x). && \text{(Zero)}
 \end{aligned}$$

1047 Let $x \in X$ and $\alpha, \alpha' \in \Lambda$. Then,

$$\begin{aligned}
 1048 \quad & (h \circ \omega_X)(in_{w_\alpha}(\alpha', x)) \\
 1049 \quad &= w_{\alpha * \alpha'}^a(\beta(x)) && \text{(def. } h \text{ and } \omega) \\
 1050 \quad &= w_\alpha^a(w_{\alpha'}^a(\beta(x))) && \text{(Mult)} \\
 1051 \quad &= w_\alpha^a(h(\alpha', x)) && \text{(def. } h) \\
 1052 \quad &= (a \circ \Sigma_{\mathcal{W}} h)(in_{w_\alpha}(\alpha', x)). && \text{(def. } w_\alpha^a \text{ and } \Sigma_{\mathcal{W}})
 \end{aligned}$$

1054 Thus, h is a $\Sigma_{\mathcal{W}}$ -homomorphism.

1055 It remains to prove uniqueness of h . Notice first that, for any $\alpha \in \Lambda$ and $x \in X$,
 1056 $(\alpha, x) = w_\alpha^\omega(\tau(x))$, where $w_\alpha^\omega = \omega_X \circ in_{w_\alpha}$ denotes the interpretation of $w_\alpha: 1 \in \Sigma_{\mathcal{W}}$ in
 1057 $(\Lambda \square X, \omega_X)$. Indeed, the following holds

$$\begin{aligned}
 1058 \quad & (\alpha, x) = (\alpha * 0, x) && \text{(0 identity)} \\
 1059 \quad &= w_\alpha^\omega(0, x) && \text{(def. } \omega) \\
 1060 \quad &= w_\alpha^\omega(\tau(x)). && \text{(def. } \tau)
 \end{aligned}$$

1062 Assume there exists $g: \Lambda \square X \rightarrow A$ such that $g \circ \tau_X = \beta$ and $g \circ \omega_X = a \circ \Sigma_{\mathcal{W}}g$. Then, the
 1063 following holds:

$$\begin{aligned}
 1064 \quad h((\alpha, x)) &= h(\mathbf{w}_\alpha^\omega(\tau(x))) \\
 1065 \quad &= \mathbf{w}_\alpha^a(h(\tau(x))) && (h \text{ homo}) \\
 1066 \quad &= \mathbf{w}_\alpha^a(\beta(x)) && (h \circ \tau = \beta) \\
 1067 \quad &= \mathbf{w}_\alpha^a(g(\tau(x))) && (g \circ \tau = \beta) \\
 1068 \quad &= g(\mathbf{w}_\alpha^\omega(\tau(x))) && (g \text{ homo}) \\
 1069 \quad &= g((\alpha, x)) \\
 1070
 \end{aligned}$$

1071 Therefore, $h = g$. ◀

1072 **Proof.** (of Theorem 26) By Theorem 41, the functors $(\Lambda \square -)$ and $T_{\mathcal{W}}$ are isomorphic and the
 1073 units of the two monads coincide (up-to iso). We are left to prove that also the multiplications
 1074 coincide (up-to iso). By Theorem 41, this follows by showing that the following diagram
 1075 commutes

$$\begin{array}{ccc}
 (\Lambda \square X) & \xrightarrow{\tau_{\Lambda \square X}} & (\Lambda \square (\Lambda \square X)) & \xleftarrow{\omega_{\Lambda \square X}} & \Sigma_{\mathcal{W}}(\Lambda \square (\Lambda \square X)) \\
 & \searrow \text{id} & \downarrow \varsigma_X & & \downarrow \Sigma_{\mathcal{W}}\varsigma_X \\
 & & \Lambda \square X & \xleftarrow{\omega_X} & \Sigma_{\mathcal{W}}(\Lambda \square X)
 \end{array}$$

1077 $\varsigma_X \circ \tau_X = \text{id}$ holds since $(\Lambda \square -)$ is a monad. The right square diagram commutes as shown
 1078 below

$$\begin{aligned}
 1079 \quad &(\varsigma_X \circ \omega_{\Lambda \square X})(\text{in}_{\mathbf{w}_\alpha}(\alpha', (\alpha'', x))) \\
 1080 \quad &= \varsigma_X((\alpha * \alpha', (\alpha'', x))) && (\text{def. } \omega) \\
 1081 \quad &= (\alpha * \alpha' * \alpha'', x) && (\text{def. } \varsigma) \\
 1082 \quad &= \omega_X(\text{in}_{\mathbf{w}_\alpha}(\alpha' * \alpha'', x)) && (\text{def. } \omega_X) \\
 1083 \quad &= \omega_X(\text{in}_{\mathbf{w}_\alpha}(\varsigma_X(\alpha', (\alpha'', x)))) && (\text{def. } \varsigma) \\
 1084 \quad &= (\omega_X \circ \Sigma_{\mathcal{W}}\varsigma_X)(\text{in}_{\mathbf{w}_\alpha}(\alpha', (\alpha'', x))) && (\text{def. } \Sigma_{\mathcal{W}}) \\
 1085
 \end{aligned}$$

1086 for arbitrary $x \in X$ and $\alpha, \alpha', \alpha'' \in \Lambda$. ◀

1087 **Proof.** (of Lemma 31) Similar to [1, Lemma 8.4]. ◀